



**HAL**  
open science

# Interactions between phytovirus and aphids vectors can be modulated depending on vector transmission efficiency

Quentin Chesnais, Aude Couty, Marilyne Uzest, Véronique Brault, Ameline Arnaud

## ► To cite this version:

Quentin Chesnais, Aude Couty, Marilyne Uzest, Véronique Brault, Ameline Arnaud. Interactions between phytovirus and aphids vectors can be modulated depending on vector transmission efficiency. 18. Colloque de la Biologie de l’Insecte (CBI 2016), Jun 2016, Tours, France. , 2016, Programme CBI 2016. hal-02743178

**HAL Id: hal-02743178**

**<https://hal.inrae.fr/hal-02743178v1>**

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L’archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d’enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Poster | 7

## Interactions between phytovirus and aphids vectors can be modulated depending on vector transmission efficiency.

Quentin Chesnais<sup>1</sup>, Aude Couty<sup>1</sup>, Maryline Uzest<sup>2,3</sup>, Véronique Brault<sup>4</sup>, Arnaud Ameline<sup>1</sup>

<sup>1</sup>Ecologie et Dynamique des Systèmes Anthropisés (EDYSAN), Université Picardie Jules Verne, Amiens, France. <sup>2</sup>Centre de Coopération Internationale en Recherche Agronomique pour le Développement (CIRAD), Paris, France. <sup>3</sup>Institut national de la recherche agronomique (INRA), UMR BGPI, Montpellier, France. <sup>4</sup>Santé de la Vigne et Qualité du Vin (SVQV - INRA-UDS), Institut national de la recherche agronomique (INRA) : UMR1131, Colmar Cedex, France.

Vector-borne viruses can induce changes in the phenotype of their host plants that may in turn influence the frequency and nature of host–vector interactions. Manipulation of vector behaviour by phytoviruses usually facilitates their propagation and seems to converge according to the virus transmission mode. However, to our knowledge, this convergence has never been related to the vector transmission efficiency. We investigated the effects of infection of *Camelina sativa* by the persistent *Turnip Yellow Virus* and the semi-persistent *Cauliflower Mosaic Virus* on (i) arrestment and dispersal, (ii) feeding behaviour (using electropenetrography) and (iii) physiology of two aphid species: the polyphagous *Myzus persicae* and the Brassicaceae specialist *Brevicoryne brassicae*. Results showed TuYV-infected camelina induced positive effects on the feeding behaviour and physiology of *Myzus persicae*, a vector with high transmission efficiency; but induced negative effects on *Brevicoryne brassicae*, a poor vector of TuYV. CaMV-infected plants had similar effects on both aphid species showing equivalent transmission efficiency for this virus. In conclusion, viral manipulation of vector behaviour is likely to depend not only on the virus mode of transmission, which is generally considered as a common assumption, but also on the aphid transmission efficiency.