



HAL
open science

Cryopreservation of Pacific Oyster larvae: a limited but promising survival!

Marc Suquet, Pierrick Haffray, Anastasia Bestin, F. Malo, Catherine Labbé

► **To cite this version:**

Marc Suquet, Pierrick Haffray, Anastasia Bestin, F. Malo, Catherine Labbé. Cryopreservation of Pacific Oyster larvae: a limited but promising survival!. CRB-Anim Biennial Seminar 2015, Centre de Ressources Biologiques pour la Génomique des Animaux d'Élevages et d'Intérêt Économique. FRA., Feb 2015, Paris, France. 22 p. hal-02743321

HAL Id: hal-02743321

<https://hal.inrae.fr/hal-02743321>

Submitted on 3 Jun 2020


HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



2015

International Seminar



9 & 10 Feb 2015

MGEN - Paris



Labogena



2) Cryopreservation of Pacific Oyster larvae: a limited but promising survival!

M. Suquet¹, P. Haffray², A. Bestin, F. Malo¹ and C. Labbé³

In aquatic animals, while sperm cryopreservation methods are well established, the preservation of larvae is still far from being achieved. Compared to fish species, mollusk oocytes are generally small and have a lower yolk content. This is favorable to larval cryopreservation. Because of the soaring role of hatcheries, Pacific oyster farming requires larval cryopreservation.

Larvae were cryopreserved using the following conditions (Microdigitcool, IMV): larval dilution 1V:1V in cryoprotectant (10% ethylene glycol, 1% PVP, 200mM sucrose), 500µl straws, -35°C after 1h30 and then liquid nitrogen, thawing at 37°C. Survival was similar when larvae were cryopreserved at trochophore (13hpf: hours post fertilization) or D-larval stage (24hpf). A higher larval survival was observed when shortening the cryopreservation cycle from 1h30 to 45min. After a 3 year rearing period, the growth and reproductive performances of oysters, formerly cryopreserved at larval stages, were similar to those observed for unfrozen ones.

The survival yields of thawed larvae (close to 0.5%) can be largely improved in the future but the high quality of thawed larvae shows that this technique can be applicable for the establishment of Pacific oyster larvae cryobanks.

¹Ifremer, UMR 6539, PFOM Department, Argenton

²INRA, UR 1037, LPGP, Rennes

³Sysaaf, LPGP, Rennes

Acknowledgments: funded by CRECHE (Ofimer 136-08-C) and CRB-Anim (ANR-11-INBS-0003).

Marc Suquet
IFREMER

Marc Suquet is researcher in Ifremer (Center of Brest). He has a 20 year experience on fish reproduction and mainly in gamete biology and management. Since 2006, He is involved in the same subjects but now in mollusks and mainly Pacific oyster and scallop.