



## **Weight status and alcohol intake modify the association between vitamin D and breast cancer risk**

Melanie Deschasaux, Jean-Claude Souberbielle, Paule Latino Martel, Angela Sutton, Nathalie Charnaux, Nathalie Druesne Pecollo, Pilar Galan, Serge Hercberg, Sigrid Le Clerc, Emmanuelle Kesse-Guyot, et al.

### **► To cite this version:**

Melanie Deschasaux, Jean-Claude Souberbielle, Paule Latino Martel, Angela Sutton, Nathalie Charnaux, et al.. Weight status and alcohol intake modify the association between vitamin D and breast cancer risk. IARC's 50th Anniversary, Jun 2016, Lyon, France. , 518 p., 2016, IARC's 50th Anniversary "Cancer: Occurrence, Causes, Prevention". Abstracts. hal-02743355

**HAL Id: hal-02743355**

**<https://hal.inrae.fr/hal-02743355>**

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## **B-067 - Weight Status And Alcohol Intake Modify The Association Between Vitamin D And Breast Cancer Risk**

MÉLANIE DESCHASAU, SORBONNE PARIS CITÉ EPIDEMIOLOGY AND STATISTICS RESEARCH CENTER (CRESS), INSERM U1153, INRA U1125, CNAM, PARIS 13 UNIVERSITY, FRANCE

SOUBERBIELE J.<sup>2</sup>, LATINO-MARTEL P.<sup>1</sup>, SUTTON A.<sup>3</sup>, CHARNAUX N.<sup>3</sup>, DRUESNE-PECOLLO N.<sup>1</sup>, GALAN P.<sup>1</sup>, HERCBERG S.<sup>1,4</sup>, LE CLERC S.<sup>5</sup>, KESSE-GUYOT E.<sup>1</sup>, EZZEDINE K.<sup>6</sup>, TOUVIER M.<sup>1</sup>

<sup>1</sup> Sorbonne Paris Cité Epidemiology and Statistics Research Center (CRESS), Inserm U1153, Inra U1125, Cnam, Paris 13 University, Nutritional Epidemiology Research Team (EREN), 93017 Bobigny, France

<sup>2</sup> Physiology Department, Necker Hospital, Inserm U845, 75015 Paris, France

<sup>3</sup> Biochemistry Department, Jean Verdier Hospital, Inserm U698, Paris 13 University, 93140 Bondy, France

<sup>4</sup> Public Health Department, Avicenne Hospital, 93017 Bobigny, France

<sup>5</sup> Conservatoire National des Arts et Métiers (CNAM), Genomics, Bioinformatics and Applications Team (EA4627), 75003 Paris, France

<sup>6</sup> Dermatology Department, Saint André Hospital, 33000 Bordeaux, France

### **Purpose:**

Mechanistic hypotheses suggest that vitamin D may contribute to the prevention of breast cancer. However, epidemiologic evidence is inconsistent, suggesting a potential effect modification by individual factors. Our objective was to investigate the prospective associations between the plasma 25-hydroxyvitamin D (25(OH)D) concentration, polymorphisms of genes encoding for the vitamin D receptor (VDR), and the vitamin D-binding protein (GC), and breast cancer risk, along with 2 potential modifiers: BMI and alcohol intake.

### **Methods:**

A nested case-control study was set up in the Supplémentation en Vitamines et Minéraux Antioxydants (SU.VI.MAX) cohort (1994–2007), involving 233 women with breast cancer and 466 matched controls. The plasma total 25(OH)D concentration and gene polymorphisms were assessed on samples obtained at baseline. Conditional logistic regression models were computed.

### **Results:**

A higher plasma 25(OH)D concentration was associated with a decreased risk of breast cancer for women with a BMI<sub>Q4vs.Q1</sub>=0.46 (0.23-0.89); P-trend=0.01, P-interaction=0.002, whereas it was associated with an increased risk in women with a BMI ≥median (OR<sub>Q4vs.Q1</sub>=2.45 (1.13-5.28); P-trend=0.02, P-interaction=0.002). A plasma 25(OH)D concentration ≥10ng/mL was associated with a decreased risk of breast cancer for women with alcohol intakes ≥median (OR<sub>≥10vs.<10ng/mL</sub>=0.50 (0.26-0.95); P=0.03, P-interaction=0.03). The genetic analyses were consistent with the results observed with plasma 25(OH)D.

### **Conclusions:**

In this prospective study, plasma 25(OH)D was associated with a decreased breast cancer risk in lean women, whereas it was associated with an increased risk in women with a higher BMI. Plasma 25(OH)D was also associated with a decreased breast cancer risk in women with moderate-to-high alcohol consumption, whereas no association was observed among women with low alcohol intake. These effect modifications suggest explanations for discrepancies in results of previous studies.

J Nutr 2016

### **Funding source:**

French Research Institute for Public Health (IRESP), grant number AAR201206  
Cancéropôle Ile-de-France, Ile-de-France Region, PhD grant (Mélanie Deschasaux)