



**HAL**  
open science

## The diversity of the N<sub>2</sub>O reducers matters for the relative proportion of N<sub>2</sub>O emitted across cropping systems by denitrification

Luiz Domeignoz Horta, Aymé Spor, David Bru, Marie-Christine Breuil, Florian Bizouard, Joël J. Léonard, Laurent Philippot

### ► To cite this version:

Luiz Domeignoz Horta, Aymé Spor, David Bru, Marie-Christine Breuil, Florian Bizouard, et al.. The diversity of the N<sub>2</sub>O reducers matters for the relative proportion of N<sub>2</sub>O emitted across cropping systems by denitrification. 20th European Nitrogen Cycle Meeting, ENC2015 Conference, University of Aberdeen. GBR., Sep 2015, Aberdeen, United Kingdom. 65 p. hal-02743401

**HAL Id: hal-02743401**

**<https://hal.inrae.fr/hal-02743401>**

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# ORAL PRESENTATIONS

# Diversity of N<sub>2</sub>O reducers matters for the relative proportion of N<sub>2</sub>O emitted across cropping systems by denitrification

Luiz Domeignoz-Horta, A. Spor, D. Bru, M.C. Breuil, F. Bizouard, J. Léonard, Laurent Philippot

INRA, UMR 1347 Agroécologie, F-21065 Dijon cedex, France

[luizdomeignoz@gmail.com](mailto:luizdomeignoz@gmail.com)

Agriculture is the main source of terrestrial emissions of N<sub>2</sub>O, a potent greenhouse gas and the main cause of ozone layer depletion. The reduction of N<sub>2</sub>O into N<sub>2</sub> by microorganisms carrying the nitrous oxide reductase gene (*nosZ*) is the only biological process known to eliminate this greenhouse gas. Recent studies showed that a previously unknown clade of N<sub>2</sub>O-reducers was related to the capacity of the soil to act as an N<sub>2</sub>O sink, opening the way for new strategies to mitigate emissions. Here, we investigated whether the agricultural practices could differently influence the two N<sub>2</sub>O reducer clades with consequences for denitrification end-products. The abundance of N<sub>2</sub>O-reducers and producers was quantified by real-time PCR, and the diversity of both *nosZ* clades was determined by 454 pyrosequencing. Potential N<sub>2</sub>O emissions and potential denitrification activity were used to calculate the N<sub>2</sub>O emission ratio. Overall, the results showed limited differences between management practices but there were significant differences between cropping systems in both the abundance and structure of the *nosZII* community, as well as in the N<sub>2</sub>O emission ratio. More limited differences were observed in the *nosZI* community, suggesting that the newly identified *nosZII* clade is more sensitive than *nosZI* to environmental changes. Potential denitrification activity and potential N<sub>2</sub>O emissions were explained mainly by the soil properties while the diversity of the *nosZII* clade on its own explained 26 percent of the proportion of N<sub>2</sub>O emitted, which highlights the importance of understanding the ecology of this newly identified clade of N<sub>2</sub>O reducers for mitigation strategies.