



**HAL**  
open science

## Evidence for foliar uptake of sulfated laminarin into grapevine depending on surfactant use and leaf surface

Yuko Krzyżaniak, Franck Paris, Christian Gauvrit, G. Lecollinet, Vincent Ferrières, Marielle Adrian, Laurent Legentil, Xavier Daire, Sophie Trouvelot

### ► To cite this version:

Yuko Krzyżaniak, Franck Paris, Christian Gauvrit, G. Lecollinet, Vincent Ferrières, et al.. Evidence for foliar uptake of sulfated laminarin into grapevine depending on surfactant use and leaf surface. 11. International Conference on Plant Diseases, Association Française de Protection des Plantes (AFPP). FRA., Dec 2015, Tours, France. hal-02743463

**HAL Id: hal-02743463**

**<https://hal.inrae.fr/hal-02743463>**

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## **Promoting an evolutionary approach of herbicide resistance.**

Christophe Délye \*, Valérie Le Corre \*, R<sub>4</sub>P #

\* *INRA, UMR1347 Agroécologie, 17 rue de Sully, F-21000 Dijon, France*

# *R<sub>4</sub>P consortium, r4p@listes.inra.fr*

E-mail: [delye@dijon.inra.fr](mailto:delye@dijon.inra.fr)

Resistance rapidly and continuously evolves in weed populations undergoing herbicide selective pressure, so that the long-term outcome of herbicide selection remains unpredictable. Most published herbicide resistance studies focussed on the resistance mechanisms, with diagnosis as an applied aim. While elucidating resistance genetic bases is a prerequisite to understand and predict resistance evolution, resistance studies should go beyond the mere description of mechanisms. This means implementing an evolutionary approach of resistance by investigating the probability of appearance or initial frequency of resistance genes, the genetic architecture of resistance, resistance spread and the associated pleiotropic effects (e.g., fitness cost) of resistance mechanisms. This data is necessary to design integrated weed management strategies reducing herbicide resistance risk or enabling resistance management *via* integrative modelling efforts. Yet, the evolutionary dynamics of herbicide resistance has rarely been fully addressed considering all these aspects. This talks aims at promoting interest for integrative evolutionary approaches of herbicide resistance. The main parameters driving herbicide resistance evolution in weeds and their respective importance in resistance evolution will be reviewed, taking into account weed biological particularities and using comparison with the other major pest classes (fungi and insects) to highlight particularities in herbicide resistance evolution.