



HAL
open science

Understanding the effect of alkaline pretreatments of lignocellulosic biomass on cell wall structure

Hélène Thomas, Jean Luc Verdeil, Hélène Carrère

► **To cite this version:**

Hélène Thomas, Jean Luc Verdeil, Hélène Carrère. Understanding the effect of alkaline pretreatments of lignocellulosic biomass on cell wall structure. 12. International Conference on Renewable Resources and Biorefineries (RRB-12)), Ghent University [Belgium] (UGENT). BEL., May 2016, Ghent, Belgium. hal-02743673

HAL Id: hal-02743673

<https://hal.inrae.fr/hal-02743673>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

UNDERSTANDING THE EFFECT OF ALKALINE PRETREATMENTS OF LIGNOCELLULOSIC BIOMASS ON CELL WALL STRUCTURE

Hélène Laurence Thomas¹, J-L Verdeil², H Carrère¹

¹INRA, UR0050, Laboratoire de Biotechnologies de l'Environnement (LBE), Narbonne, France

²CIRAD, UMR AGAP, Hystocytology and Plant Cell Imaging Platform, Montpellier, France

helene.thomas@supagro.inra.fr

The emerging biorefinery concept is highly attractive to optimize lignocellulosic biomass use (Ghatak 2011). Sorghum has a high genotypic and phenotypic variability and it presents several environmental advantages as it needs few inputs. This crop has a high aboveground biomass production with a high valorization potential and can also be grown as catch crop. It has been studied and used for several energy vectors production such as bioethanol, biohydrogen or biogas. All the biological processes rely on cellulose and hemicelluloses hydrolysis and fermentation; however polysaccharides accessibility in lignocellulosic biomass is limited by lignin and its links with polysaccharides. Consequently, biomass pretreatment is a key step in biorefinery for cell wall deconstruction and increase the accessibility of sugars to enzymes and microorganisms. Among efficient pre-treatments for delignification, chemicals and more precisely alkaline have been highlighted prior both enzymatic hydrolysis in bioethanol process (Mood et al., 2013) and anaerobic digestion (Monlau et al. 2012).

The aim of this study was to understand and compare the mechanisms of alkaline pretreatments on sorghum stem cell wall. For this purpose, three pretreatments (NaOH 10g/100g_{TS}, CaO 10g/100g_{TS} and NH₄OH 1L/100g_{TS}) in liquid medium (90 g_{TS}/L) were applied on 1 mm stem fragments of two genotypes (B140 and IS 30405). To analyse the impact of alkali pretreatment on sorghum cell wall structure, we conducted an original histochemical study on raw and pretreated biomass using FASGA staining (Tolivia and Tolivia 1987) and phloroglucinol lignin staining. In this way, we monitored the plant cell wall change. In addition cellulose, hemicelluloses and lignin contents were measured by acid hydrolysis (Table 1).

Table 1: Composition of raw and pretreated sorghum genotypes

	Hemicelluloses (%TS)	Cellulose (%TS)	Klason lignin (% TS)
B 140	18.3 ± 0.4	27.7 ± 0.8	13.7 ± 1.5
B140 NaOH	15.4 ± 1	23.1 ± 1.8	13.3 ± 1.1
B140 CaO	14.8 ± 0.2	20.8 ± 1	12.1 ± 0.9
B 140 NH ₄ OH	15.4 ± 2.5	24.9 ± 1.9	nd
IS 30405	20.4 ± 0.4	31.6 ± 0.3	17.8 ± 0.2
IS NaOH	18.8 ± 0.06	29.3 ± 0.1	16.9 ± 0.6
IS CaO	18.0 ± 0.2	26.9 ± 1.1	16.3 ± 0.8
IS NH ₄ OH	11.8 ± 1.9	19.3 ± 1.2	nd

References

- Ghatak, Himadri Roy. 2011. "Biorefineries from the Perspective of Sustainability: Feedstocks, Products, and Processes." *Renewable and Sustainable Energy Reviews* 15 (8): 4042–52.
- Monlau, F., A. Barakat, J.P. Steyer, and H. Carrere. 2012. "Comparison of Seven Types of Thermo-Chemical Pretreatments on the Structural Features and Anaerobic Digestion of Sunflower Stalks." *Bioresource Technology* 120 (September): 241–47.
- Mood SH, Golfeshan AH, Tabatabaei M, Jouzani GS, Najafi G, Gholami M, Ardjmand M. Lignocellulosic biomass to bioethanol, a comprehensive review with a focus on pretreatment. *Renewable & Sustainable Energy Reviews*. 2013;27:77-93
- Tolivia, D., and J. Tolivia. 1987. "Fasga: New Polychromatic Method for Simultaneous and Differential Staining of Plant Tissues." *Journal of Microscopy*, 113–17.

Comment citer ce document :

Thomas, H. (Auteur de correspondance), Verdeil, J. L., Carrère, H. (2016). Understanding the effect of alkaline pretreatments of lignocellulosic biomass on cell wall structure. In: Final Program & Abstract Book (p. p 37). Presented at 12. International Conference on Renewable Resources and Biorefineries (RRB-12). Ghent, BEL (2016-05-30 - 2016-06-01).