



HAL
open science

Cryopreserving European eel (*A. anguilla*) sperm: comparison of two methods for standardization

Maria C. Vilchez, Marina Morini, David S. Penaranda, Luz Pérez, Alexandra Depince, Esztzer Kasa, Catherine Labbé, Akos Horvath, Juan F. Asturiano

► **To cite this version:**

Maria C. Vilchez, Marina Morini, David S. Penaranda, Luz Pérez, Alexandra Depince, et al.. Cryopreserving European eel (*A. anguilla*) sperm: comparison of two methods for standardization. 5. Jornadas Ibéricas de Ictiología - SIBIC2014, Jun 2014, Lisboa, Portugal. hal-02743772

HAL Id: hal-02743772

<https://hal.inrae.fr/hal-02743772>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Tipo de comunicación preferida: **Oral**

Cryopreserving European eel (*A. anguilla*) sperm: comparison of two methods for standardization

María C. VÍLCHEZ¹, Marina MORINI^{1,2}, David S. PEÑARANDA¹, Luz PÉREZ¹, Alexandra DEPINCE², Eszter KÁSA^{3,1}, Catherine LABBÉ², Ákos HORVATH^{3,1}, **Juan F. ASTURIANO**¹

¹ **Grupo de Acuicultura y Biodiversidad. Instituto de Ciencia y Tecnología Animal. Universitat Politècnica de València, Camino de Vera s/n. Edificio 7G. 46022 Valencia (Spain). (jfastu@dca.upv.es)**

² INRA, Fish Physiology and Genomics UR 1037, Campus de Beaulieu, Rennes France.

³ Department of Aquaculture, Szent István University, Páter Károly u. 1., H-2100 Gödöllő, Hungary

Summary

During the last years two groups of research have been working in parallel to develop protocols to cryopreserve European eel sperm. Each group established their own protocols in Spain (Asturiano et al., 2003; Peñaranda et al., 2009) and Hungary (Müller et al., 2004; Szabó et al., 2005). These methods are completely different in terms of cryoprotectant, volumes, freezing media, grade for dilution, etc., making clear the need of standardization (which is a usual situation in fish species).

Recent experiments being part of the PRO-EEL project (www.pro-eel.eu) proved the efficacy of the “Spanish method” by producing larvae of this species using previously frozen-thawed sperm (Asturiano et al., 2013), although the results should be improved because of the low fertilization rates. On the other hand, the Hungarian team was able to generate hybrid larvae using semen of European eel and Japanese eel eggs (Müller et al., 2012), evidencing the validity of the “Hungarian method”.

In the framework of the AQUAGAMETE COST Action (aquagamete.webs.upv.es), a series of joint experiments were carried out in order to standardize eel cryopreservation procedures. Males were matured with weekly injections of hCG and after 10 weeks of treatment sperm samples were extracted. Sperm motility was determined after activation with sea water, and only those having more than 68% of motile cells were selected for the experiment. Sperm



samples were frozen using both protocols. In general, sperm was diluted with a mixture of extender and cryoprotectant, loaded into straws and frozen in the vapor of liquid nitrogen. Straws were thawed by immersion into water bath for a given period of time.

Fresh and post-thawed samples were evaluated using sperm motility and morphometry parameters (using CASA and ASMA software, respectively). Moreover, because sperm cryopreservation in fish often depends on the use of permeating cryoprotectants bearing chemically reactive methyl group, we wanted to decipher whether eel sperm DNA methylation pattern was affected by the different cryopreservation protocols. The degree of DNA methylation was evaluated in fresh and frozen-thawed samples using the restriction enzyme assay and assessing methylation by image analysis and by LUMA.

The “Hungarian method” caused the higher sperm motility results post-thawing. Spermatozoa morphometry analyses and epigenetics evaluation are still in course.

Acknowledgements: Funded by EU (PRO-EEL, grant agreement n°245257) and by the CRB-Anim project "Investissements d'Avenir" ANR-11-INBS-0003. M.M., E.K. and Á.H. had grants from the COST Office (COST Action FA1205: AQUAGAMETE).