



HAL
open science

Comparison of different Marker-Assisted BLUP models for a new French genomic evaluation

Pascal Croiseau, Aurélia A. Baur, David Jonas, Chris Hoze Hoze, Julie
Promp, Didier Boichard, Sebastien S. Fritz, Vincent Ducrocq

► **To cite this version:**

Pascal Croiseau, Aurélia A. Baur, David Jonas, Chris Hoze Hoze, Julie Promp, et al.. Comparison of different Marker-Assisted BLUP models for a new French genomic evaluation. 66. Annual Meeting of the European Federation of Animal Science (EAAP), European Federation of Animal Science (EAAP). INT., Aug 2015, Warsaw, Poland. hal-02743878

HAL Id: hal-02743878

<https://hal.inrae.fr/hal-02743878v1>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Session 22 – Page 248

Comparison of different Marker-Assisted BLUP models for a new French genomic evaluation

P. Croiseau¹, A. Baur², D. Jonas^{1,2,3}, C. Hoze², J. Promp⁴, D. Boichard¹, S. Fritz², V. Ducrocq¹

¹INRA, UMR1313 GABI, 78350 Jouy-en-Josas, France

²ALLICE, 149 rue de Bercy, 75012 Paris, France

³AGROPARISTECH, 16 rue Claude Bernard, 75231 Paris, France

⁴IDELE, 149 rue de Bercy, 75012 Paris, France

French genomic evaluations have been based on a Marker-Assisted BLUP (MABLUP) approach where QTL were detected using a variable selection method. The evaluation software was modified to remove the existing constraint on number of QTL included. A set of new MABLUP models were tested using a set of 2574 Montbéliard bulls genotyped on the Illumina Bovine SNP50 BeadChip®. From 1000 to 6000 QTL detected with a BayesCpi approach were included. A model based on SNP only was compared with models based on haplotypes of 3, 4, or 5 SNP. The best SNP corresponding to a QTL position was completed with 2, 3 or 4 flanking SNP to build a haplotype. An alternative to this strategy was developed to select the haplotype with the largest number of “predictable” alleles from a window of 10 SNP around the QTL-SNP, where observed haplotype allele frequencies were used as indicators of “good predictability” in genomic evaluation. In the MABLUP model, a polygenic residual component explaining 10 to 50% of the genetic variance was also included in the model and was estimated either through the pedigree or from the SNP of the Illumina Bovine LD chip. The impact of all these scenarios on the correlation between observed daughter yield deviations and genomic estimated breeding values and on the regression slope in a regular validation test was measured and compared to other genomic selection approaches. The best combinations of these scenarios led to an average gain in correlation between 1.5 and 3% with a significant improvement of the regression slope. A reasonable final model for most traits includes 3K QTL haplotypes of 4 SNP and a residual polygenic component based on the LD SNP explaining 20 to 30% of the genetic variance. These results were used to improve the new French national genomic evaluation.