

Genotype by watering regime interaction in cultivated tomato: from phenotypes to genes

Elise Albert, Yolande Carretero, Justine Gricourt, Esther Pelpoir, Romain Novaretti, Claire Duffes, Julien Bonnefoi, Nadia Bertin, Frederique Bitton, Stephanie Pateyron, et al.

▶ To cite this version:

Elise Albert, Yolande Carretero, Justine Gricourt, Esther Pelpoir, Romain Novaretti, et al.. Genotype by watering regime interaction in cultivated tomato: from phenotypes to genes. 12. Solanaceae Conference, Oct 2015, Bordeaux, France. hal-02743885

HAL Id: hal-02743885 https://hal.inrae.fr/hal-02743885

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



The 12th Solanaceae Conference

October 25 -29, 2015 ENSEIRB Building, Talence

Bordeaux, France



Genotype by watering regime interaction in cultivated tomato: from phenotypes to genes.

Albert E^a, Carretero Y^a, Gricourt J^a, Pelpoir E^a, Novaretti R^a, Duffes C^a, Bonnefoi J^b, Bertin N^c, Bitton F^a, Pateyron S^d, Tamby JP^d, Causse M^a

In the next decade water will be limiting crop production, in particular in Mediterranean regions. Studying genotype x water regime interactions is needed to improve plant adaptation to drought. In response to environmental constraints, plants can change their phenotypes (at molecular, physiological and morphological levels). In Tomato, extensively grown in Mediterranean regions, first studies have shown genetic variability in the response to drought, but few genes/QTLs have been identified and mostly in wild related species. Studying water deficit in this crop is of particular interest since a mastered water deficit can stimulate metabolite production, increasing plant defenses and concentration of compounds involved in fruit quality at the same time. We analyzed two populations: recombinant inbreed lines (RILs) and unrelated cherry tomato accessions, grown in greenhouse under two watering regimes. We assessed a large genetic variability and highly significant genotype x water regime interactions, for several plant and fruit traits, in the two populations. Large fruit accessions showed high sensitivity to drought. The two panels were genotyped with large sets of SNP and quantitative trait loci (QTLs) were identified, combining linkage and association mapping. 20% of the QTLs were interactive between the watering regimes, mostly with antagonist effects according to treatment. Analysis whole genome of gene expression in young leaves from the RIL parents provided interesting candidate genes under interactive QTLs.

^a INRA, UR1052, Génétique et Amélioration des Fruits et Légumes, CS 60094, 84143 Montfavet, France

^b GAUTIER Semences, 13630 Eyragues, France

c INRA, UR1115, Plante et Système de cultures Horticoles, CS60094, 84143 Montfavet, France

^d INRA, Institut of Plant Sciences Paris-Saclay, UMR 9213/UMR1403, CNRS, INRA, Université Paris-Sud, Université d'Evry, Université Paris-Diderot, Sorbonne Paris-Cité, 91405 Orsay, France