



Mapping averaged pairwise information (MAPI): a new tool for landscape genetics on plant virus

Sylvain S. Piry, Marie Pierre Chapuis, Bertrand Gauffre, Julien Papaix,
Astrid Cruaud, Karine Berthier

► To cite this version:

Sylvain S. Piry, Marie Pierre Chapuis, Bertrand Gauffre, Julien Papaix, Astrid Cruaud, et al.. Mapping averaged pairwise information (MAPI): a new tool for landscape genetics on plant virus. 13. International plant virus epidemiology symposium, Jun 2016, Avignon, France. 165 p. hal-02743997

HAL Id: hal-02743997

<https://hal.inrae.fr/hal-02743997>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

MAPPING AVERAGED PAIRWISE INFORMATION (MAPI): A NEW TOOL FOR LANDSCAPE GENETICS ON PLANT VIRUS

Piry, S. (1), Chapuis, M.P. (2), Gauffre, B. (3,4), Papaïx, J. (5), Cruaud, A. (1), Berthier, K. (6)

(1) INRA, UMR 1062, Centre de Biologie pour la Gestion des Populations, 755 avenue du campus Agropolis, 34988 Montferrier sur Lez, France.

(2) CIRAD, UMR 1062, Centre de Biologie pour la Gestion des Populations, 755 avenue du campus Agropolis, 34988 Montferrier sur Lez, France.

(3) CEBC-CNRS (UPR 1934), F-79360, Beauvois sur Niort, France

(4) INRA, USC1339 (CEBC-CNRS), F-79360, Beauvois sur Niort, France

(5) INRA, UR546 Biostatistique et Processus Spatiaux, F-84914 Avignon, France

(6) INRA, UR0407 pathologie végétale, F-84143 Montfavet, France – karine.berthier@avignon.inra.fr



BACKGROUND and OBJECTIVES:

Visualisation of spatial networks based on pairwise metrics such as (dis)similarity coefficients provides direct information on spatial organization of biological systems. However, for large networks, graphical representations are often unreadable as nodes (samples), and edges (links between samples) strongly overlap. We present a new method, MAPI (Mapping Averaged Pairwise Information), allowing translation from spatial networks to variation surfaces.

MATERIALS and METHODS:

MAPI relies on i) a spatial network in which samples are linked by ellipses and, ii) a grid of hexagonal cells encompassing the study area. Pairwise metric values are attributed to ellipses and averaged within the cells they intersect. The resulting surface of variation can be displayed as a colour map in GIS, along with other relevant layers, such as land cover. The method also allows the identification of significant discontinuity in grid cell values through a nonparametric randomisation procedure.

RESULTS.

To illustrate our method we analysed *high-throughput* sequencing data of the Endive Necrotic Mosaic Virus virus obtained from 343 wild salsify collected over ~1500 km² in South Eastern France. The MAPI analysis revealed a strong spatial genetic structure that could mostly result from host plant population dynamics due to habitat heterogeneity.

CONCLUSIONS:

MAPI is freely provided as a PostgreSQL/PostGIS database extension allowing easy interaction with GIS software and other programming languages. Although developed for spatial and landscape genetics, the method can also be useful to visualise spatial organisation from other kinds of data from which pairwise metrics can be computed.

Building bridges between disciplines for sustainable management of plant virus diseases



13th International Plant Virus Epidemiology Symposium
6-10 june 2016, Avignon, FRANCE

Programme and Abstracts