



Combining various biological methods to control powdery mildew of tomato

Marc Bardin, Lea Dantony, Magali Duffaud, Laurent Neu, Michel Pascal, Claire Troulet, Philippe C. Nicot

► To cite this version:

Marc Bardin, Lea Dantony, Magali Duffaud, Laurent Neu, Michel Pascal, et al.. Combining various biological methods to control powdery mildew of tomato. 13. IOBC-WPRS Meeting of the working group "Biological control of fungal and bacterial plant pathogens". Biocontrol of plant diseases: "from the field to the laboratory and back again", Jun 2014, Uppsala, Sweden. 2014. hal-02744060

HAL Id: hal-02744060

<https://hal.inrae.fr/hal-02744060>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Combining various biological methods to control powdery mildew of tomato

Marc Bardin, Lea Dantony, Magali Duffaud, Laurent Neu, Michel Pascal, Claire Troulet, Philippe Nicot, INRA, UR407, Plant Pathology Unit, Domaine St Maurice, F-84140 Montfavet, France.
E-mail: marc.bardin@avignon.inra.fr

Various biological methods, mainly based on the utilization of antagonistic microorganisms or plant extracts, have been studied to control powdery mildews. The hyperparasite fungus *Ampelomyces quisqualis* (AQ-10, De Sangosse) is registered in many countries to control powdery mildew on various crops, including *Oidium neolycopersici* on tomato in France. A plant extract from orange (Prev-Am, Vivagro) is registered to control powdery mildew and whitefly on various crops but its effect on tomato powdery mildew is not clearly established. Spray of soluble carbohydrates on leaves of various plant species proved to have an effect against some plant pests but their effect against plant pathogens is not well known, except for trehalose against powdery mildew of wheat.

The objectives of this study were (i) to test the effect of spraying low concentration of sugars (glucose, fructose, sucrose, trehalose) on leaves of tomato against *O. neolycopersici* in controlled conditions and (2) to evaluate the effect of sucrose, AQ-10 and Prev-Am separately and in combinations in semi-commercial tunnel conditions.

Results revealed that spray of sugars on tomato leaves have no significant effect against powdery mildew in controlled conditions. In tunnels, the plant extract Prev-am showed a significant effect against *Oidium neolycopersici* on tomato. Combination of products did not enhance the efficacy of a single product. The relevance of these results to ensure the control of powdery mildew on tomato without fungicides will be discussed.



International Organisation for Biological and Integrated Control of Noxious Animals and Plants (IOBC)

West Palaearctic Regional Section (WPRS)

Organisation Internationale de Lutte Biologique et Intégrée contre les Animaux et les Plantes Nuisibles (OILB)

Section Régionale Ouest Paléarctique (SRPOP)

XIII Meeting of the Working Group Biological control of fungal and bacterial plant pathogens

Viticulture

Horticulture

Agriculture

Forestry

Biocontrol of Plant Diseases: “From the field to the laboratory and back again”



Uppsala, Sweden
SLU, Swedish University of
Agricultural Sciences



15 – 18 June 2014

Programme Abstracts Participants

