



HAL
open science

Light and temperature of ripening tomato fruits interact to regulate ascorbate synthesis, oxidation and recycling

Hélène Gautier, Capucine Massot, Doriane Bancel, Vincent Truffault, Rebecca Stevens

► To cite this version:

Hélène Gautier, Capucine Massot, Doriane Bancel, Vincent Truffault, Rebecca Stevens. Light and temperature of ripening tomato fruits interact to regulate ascorbate synthesis, oxidation and recycling. 11. International POG Conference Reactive Oxygen and Nitrogen Species in Plants, Jul 2013, Varsovie, Poland. hal-02744410

HAL Id: hal-02744410

<https://hal.inrae.fr/hal-02744410>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Conference Paper: Light and temperature of ripening tomato fruits interact to regulate ascorbate synthesis, oxidation and recycling

[H Gautier](#) · [C Massot](#) · [D Bancel](#) · [V Truffault](#) · R Stevens

[\[Show abstract\]](#)

BioTechnologia. Journal of Biotechnology Computational Biology and Bionanotechnology, Warsaw; 01/2013

Conference: BioTechnologia. Journal of Biotechnology Computational Biology and Bionanotechnology, At Warsaw, Volume: vol. 94(2) pp. 174 2013

Abstract

To understand how light and temperature may affect fruit ascorbate (AsA) content, tomato fruit harvested at breaker stage were placed under different temperatures (12, 23 and 31EC) and irradiance regimes (darkness or 150 $\mu\text{mol m}^{-2} \text{s}^{-1}$). After 56h, fruit were cut and pericarps were frozen and ground in liquid nitrogen. Changes in AsA metabolism were characterized from ascorbate and glutathione content, enzymic activities related to oxidative stress and AsA/glutathione cycle (MDHAR, DHAR, APX, CAT, GR) and the expression of genes coding for the 5 last enzymes of AsA biosynthesis pathway (GME, GDP, GPP, GalDH, GLDH). It confirms the important role of fruit micro-climate to regulate fruit AsA content and reveals interaction between light and temperature: Indeed light increased AsA content in fruit pericarp up to 67% at 12EC, but had no effect on AsA content at 31EC. At any temperature tested, light enhanced the expression of genes coding for AsA biosynthesis, but at 12EC, light upregulated a higher amount of genes compared to 23EC or 31EC. At 31EC, reductase activities (MDHAR and GR) were significantly reduced under light indicating that enzymes of the ascorbate/glutathione cycle may be limiting to recycle AsA.