



**HAL**  
open science

## **NS3 of bluetongue virus interferes with the innate antiviral response**

Damien Vitour, Estelle Lara, Emilie Chauveau, Micheline Adam, Corinne Sailleau, Emmanuel Breard, Stéphan Zientara

### ► **To cite this version:**

Damien Vitour, Estelle Lara, Emilie Chauveau, Micheline Adam, Corinne Sailleau, et al.. NS3 of bluetongue virus interferes with the innate antiviral response. 9. Joint Meeting of International Cytokine Society and International Society for Interferon and Cytokine Research, International Cytokine Society. ITA., Oct 2011, Florence, Italy. 10.1016/j.cyto.2011.07.421 . hal-02744463

**HAL Id: hal-02744463**

**<https://hal.inrae.fr/hal-02744463>**

Submitted on 6 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## **NS3 of Bluetongue virus interferes with the innate antiviral response**

**Estelle LARA, Emilie CHAUVEAU, Virginie DOCEUL, Micheline ADAM, Corinne SAILLEAU, Emmanuel BREARD, Cyril VIAROUGE, Alexandra DESPRAT, Stéphan ZIENTARA, Damien VITOUR.**

Bluetongue disease is a major animal health concern transmitted through the bites of *Culicoides* vectors. Bluetongue virus (BTV), the etiologic agent of the disease, is a dsRNA virus belonging to the genus *Orbivirus*, into the *Reoviridae* family. BTV infection triggers the production of type-I interferon (IFN-I) and genomic dsRNA is a strong IFN-I inducer. We recently showed that RIG-I-like receptor pathway is involved in the innate immune response following BTV dsRNA transfection. Most of viruses have evolved versatile strategies to escape the IFN-I response but nothing is known on the ability of BTV to counteract the innate response. In this study, we demonstrated that BTV serotype 8 (BTV-8) can dampen the type-I interferon response and that the non-structural protein 3 (NS3) is involved in this process. In order to identify viral components involved in this inhibition and to explore its possible link with viral pathogenesis, we performed a yeast-two hybrid screen using NS3 BTV-8 as bait and a human cDNA library as prey. We describe here the result of this screen and discuss its potential link to virulence strength. Amino acids sequences/residues essential for the dedicated interaction will be changed onto the corresponding gene segment in a reverse genetic system to assess whether innate response can be restored at the virus level.

Congres annuel de la Société Française d'Immunologie

8-10 NOVEMBER 2011 - MONTPELLIER