



**HAL**  
open science

## **Experimental evidence of the efficiency of two resistance genes deployment strategies - pyramiding or alternating - for sustainable management of root-knot nematodes**

Caroline Djian-Caporalino, Ariane Fazari, Nathalie Marteu, Anne-Marie Sage-Palloix, S. Risso, R. Lanza, Alain Palloix, Philippe Castagnone-Sereno

### ► **To cite this version:**

Caroline Djian-Caporalino, Ariane Fazari, Nathalie Marteu, Anne-Marie Sage-Palloix, S. Risso, et al.. Experimental evidence of the efficiency of two resistance genes deployment strategies - pyramiding or alternating - for sustainable management of root-knot nematodes. Plant resistance sustainability 2012. International conference, Oct 2012, La Colle sur Loup, France. hal-02744512

**HAL Id: hal-02744512**

**<https://hal.inrae.fr/hal-02744512v1>**

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# PLANT RESISTANCE SUSTAINABILITY

## International Conference 2012



La Colle-Sur-Loup (France)  
October 16th-19th, 2012



## Scientific Programme and Abstracts

### Sessions

**Session 1:** Impact of plant disease resistance on the structure and evolution of pathogen populations

**Session 3:** From plant-pathogen molecular interactions to the durability of resistance

**Session 2:** Sustainable and integrated breeding and deployment of genetic resistance

**Session 4:** Socio-economic issues related to the use of resistant varieties and their deployment in agro-systems

### Invited Speakers

**Philippe Baret**, Université Catholique de Louvain, Belgium - **James Brown**, John Innes Centre, England - **Marion Desquilbet**, INRA, France - **Sylvain Gandon**, CNRS, France - **Benoit Moury**, INRA, France - **Chris Mundt**, Oregon State University, USA - **Laura Rose**, Heinrich-Heine University, Germany - **Walter Rossing**, Wageningen University, The Netherlands - **Peter Thrall**, CSIRO Plant Industry, Australia

Organised by the Institut National de la Recherche Agronomique (INRA)  
Metaprogramme on Sustainable Management of Crop Health (SMaCH)





**Experimental evidence of the efficiency of two resistance genes deployment strategies - pyramiding or alternating - for sustainable management of root-knot nematodes**

**Djian-Caporalino C.<sup>1</sup>; Fazari A.<sup>1</sup>; Marteu N.<sup>1</sup>; Sage-Palloix A.M.<sup>2</sup>; Risso S.<sup>3</sup>; Lanza R.<sup>3</sup>; Palloix A.<sup>2</sup>; Castagnone-Sereno P.<sup>1</sup>**

<sup>1</sup>INRA PACA (Provence Alpes Côte d'Azur), UMR1355 INRA/UNSA/CNRS, Institut Sophia Agrobiotech, BP167, F-06903 Sophia Antipolis, France

<sup>2</sup>INRA PACA, UR1052, Génétique et Amélioration des Fruits et Légumes, F-84143 Montfavet, France

<sup>3</sup>Chambre d'agriculture des Alpes Maritimes, 1 route de Grenoble, 06200 Nice, France

**Abstract**

The current restrictions of the use of chemical nematicides have contributed to increase root-knot nematode (RKN) problems in horticultural crops. In this context, plant resistance appears as the most effective and sustainable method of control. In horticultural crops, a few major resistance (*R*) genes are available, but the possible occurrence of virulent nematode populations able to reproduce on *R*-plants may constitute a severe threat to this control strategy. Here, we tested several *R*-gene deployment strategies in the pepper-nematode interaction to implement a rational management of the *R*-cultivars increasing the sustainable management of RKN. Experiments were conducted in climate-controlled rooms, in greenhouses, and under 3-years-field agronomic conditions comparing i) the succession of the same *R*-gene every year, when introgressed in a resistant vs. a susceptible genetic background, ii) the alternance of single *R*-genes in rotation, iii) the mixture of genotypes bearing single *R*-genes sown in the same plot, and iv) the pyramiding of two *R*-genes in one genotype. Results allow the identification of conditions lowering the emergence of virulent nematodes and assess the time required for the sustainable improvement of soil health (reduction of parasite populations under their damage threshold) using the *R*-plants as RKN "traps". Alternating different *R*-genes in rotation was confirmed to be efficient to decrease virulent populations in fields due to the specificity of the virulence previously demonstrated in laboratory experiments. Suppression of the emergence of virulent isolates because of the pyramiding of two different *R*-genes in one pepper genotype was confirmed in controlled and natural conditions and proved durable during a 3-years-field experiment. Recently, a new project called 'GEDUNEM' has been launched in the framework of the INRA metaprogramme SMaCH (Sustainable Management of Crop Health). It aims at identifying innovative strategies across the agrosystem, combining varietal resistances in crop rotations with agronomic practices, for the sustainable and integrated management of RKN in protected vegetable cropping systems.

Keywords: sustainable crop protection, breeding strategy, resistance gene deployment, virulence emergence, root knot nematodes, *Meloidogyne spp.*, *Capsicum spp.*