

New resistance-genes deployment strategies as non chemical alternatives for the durable management of root-knot nematodes in vegetable crops rotation

Caroline Djian-Caporalino, Alain Palloix, Ariane Fazari, Nathalie Marteu, Arnaud Barbary, Anne-Marie Sage-Palloix, Sabine Risso, Roger Lanza, Catherine Taussig, Philippe Castagnone-Sereno

► **To cite this version:**

Caroline Djian-Caporalino, Alain Palloix, Ariane Fazari, Nathalie Marteu, Arnaud Barbary, et al.. New resistance-genes deployment strategies as non chemical alternatives for the durable management of root-knot nematodes in vegetable crops rotation. 2nd international symposium on organic greenhouse horticulture, International Society for Horticultural Science (ISHS). INT., Oct 2013, Avignon, France. 310 p. hal-02744854

HAL Id: hal-02744854

<https://hal.inrae.fr/hal-02744854>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Session Pest control

Preferred Oral presentation

New resistance-genes deployment strategies as non chemical alternatives for the durable management of root-knot nematodes in vegetable crops rotation.

Caroline Djian-Caporalino¹, Alain Palloix², Ariane Fazari¹, Nathalie Marteu¹, Arnaud Barbary¹, Anne-Marie Sage-Palloix², Sabine Risso⁴, Roger Lanza⁴, Catherine Taussig⁵, Philippe Castagnone-Sereno¹

¹INRA PACA (Provence Alpes Côte d'Azur), UMR1355 INRA/UNSA/CNRS, Institut Sophia Agrobiotech, F-06903 Sophia Antipolis, France <caroline.caporalino@sophia.inra.fr>

²INRA PACA, UR1052, Génétique et Amélioration des Fruits et Légumes, [CS 60094](#), F-~~84143~~ [84140](#) Montfavet, France

³Chambre d'agriculture des Alpes Maritimes, 1 route de Grenoble, F-06200 Nice, France

⁴APREL, Association Provençale de Recherche et d'Expérimentation Légumière, Route de Mollégès, 13210 Saint-Rémy de Provence, France

The current restrictions on the use of chemical nematicides have contributed to increase root-knot nematode problems in horticultural crops. In this context, plant resistance appears as the most effective method of control, but the possible occurrence of virulent nematodes able to reproduce on *R*-plants may constitute a severe threat to this control strategy. To implement a rational management of *R*-lines increasing the durability of the *R*-genes, we tested several *R*-gene deployment strategies. Experiments were conducted in climate-controlled rooms, in greenhouses, and under 3-years-field agronomic conditions comparing the succession of the same *R*-genes every year, when introgressed in a partially resistant vs. a susceptible genetic background, the alternance, the mixt cultivation and the pyramiding of two *R*-genes with different modes of action in one-a single genotype. At the plant level, we previously showed that i) the choice of the *R*-genes and the genetic backgrounds in which they are introgressed can lower the frequency of resistance breakdown, and ii) the pyramiding of two different *R*-genes in one genotype totally suppressed the emergence of virulent isolates. Here, at the field and rotation level, we confirmed these results and showed that i/ alternating different *R*-genes in rotation was efficient to reduce the selection pressure of *R*-genes on the pathogens and allowed to recycle ineffective *R*-genes, and ii) optimal cultivation practices of *R*-plants increase their "trap" effect and may decrease the amount of pathogens in the soil, below their damage threshold. These results are in good agreement with concepts recently developed from the analysis of other plant-pathogen interactions. The root-knot nematode model could thus contribute to generalize strategies for the breeding and management of *R*-cultivars strengthening and increasing the durability of qualitative resistances.

This research was carried out with the financial support from the French Groupement d'Intérêt Scientifique PICléq ('Néoleg2') and the National Research Agency ('Sysbiotel').

Keywords: Sustainable crop protection, breeding strategy, resistance gene deployment, virulence emergence, root knot nematodes, *Meloidogyne* spp., *Capsicum* spp.