



Influence of post partum nutritional level on estrus behavior in primiparous Charolais cows

Emilie Recoules, Anne de La Torre, Jacques Agabriel, David Egal, Fabienne Blanc-Vazeille

► To cite this version:

Emilie Recoules, Anne de La Torre, Jacques Agabriel, David Egal, Fabienne Blanc-Vazeille. Influence of post partum nutritional level on estrus behavior in primiparous Charolais cows. 62. Annual Meeting of the European Association for Animal Production (EAAP), Aug 2011, Stavanger, Norway. hal-02744905

HAL Id: hal-02744905

<https://hal.inrae.fr/hal-02744905>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Influence of postpartum nutritional level on estrus behavior in primiparous Charolais cows.

Recoules E.^{1,4}, De La Torre A.¹, Agabriel J.¹, Egal D.² and Blanc F^{3,4}.

¹INRA–URH1213, Theix F-63122 Saint-Genès-Champanelle, ²INRA-UE 1296, 63820 Laqueuille, ³Clermont Université, VetAgro Sup, UR 2008.03.102, EPR, BP 10448, F-63000 Clermont-Ferrand, ⁴INRA, USC 2005, F-63370, Lempdes

In mountain areas, beef cattle feeding systems mainly rely on forage resources. As feed stocks may be inadequate due to environmental constraints, animals may be underfed during winter periods. Under nutrition could affect female's reproductive performances and so economic viability of suckling farms. Estrus expression is a key component of reproductive performance, especially when artificial insemination is used. Two experiments (trial 1: n=14 and trial 2: n=16) were successively carried out using primiparous Charolais cows. Body condition scores at calving were 2.4 ± 0.13 in trial 1 and 2 ± 0.20 in trial 2 (scale 0-5). In each trial, two energy level diets (High vs Low) were applied from calving to turn out. Weight and body condition were regularly measured during the experiment. Cows were reared in groups (n= 7 or 8) in a loose-housing system from mid November to mid-May. Weight and body condition were regularly measured during the experiment. Reproductive efficiency was assessed through physiological and behavioral variables. Cyclicity was studied from plasmatic progesterone profiles. Estrus expression was analyzed from 24h/24h video camera records that were studied using The Observer® software. On average during estrus periods, social and sexual behaviors represent respectively $46\% \pm 14$, $54\% \pm 14$ of total interactions. Standing to be mounted represents only $3.7\% \pm 3.5$ of total interactions. Mean estrus duration (time between first and last standing to be mounted) is $7h \pm 6$. Mean estrus intensity, defined as the sum of all sexual behaviors during estrus period is 242 ± 157 . Mean duration of estrus is longer ($9h \pm 5$ vs $5h \pm 6$, $p < 0.05$) and intensity of estrus expression is higher (303 ± 182 vs 184 ± 100 , $p < 0.05$) in L than in H cows. This result might be explained by less time spent interacting with fellows in H cows as they spend more time eating than L cows.