



**HAL**  
open science

# Effect of nitrogen fertilisation of strawberry plants on the efficacy of defence-stimulating biocontrol products against *Botrytis cinerea*

Philippe C. Nicot, Marc Bardin, François Lecompte

► **To cite this version:**

Philippe C. Nicot, Marc Bardin, François Lecompte. Effect of nitrogen fertilisation of strawberry plants on the efficacy of defence-stimulating biocontrol products against *Botrytis cinerea*. PRIR 2011, Sep 2011, Neuchatel, Switzerland. hal-02744988

**HAL Id: hal-02744988**

**<https://hal.inrae.fr/hal-02744988v1>**

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# PR-Proteins and Induced Resistance Against Pathogens and Insects

*molecular biology meets application*



Neuchâtel, Switzerland



4-8 September 2011

## Abstract | book



PR-proteins & Induced Resistance

# PRIR+ 2011

September 4-8, Neuchâtel, Switzerland

Resistance against pathogens & insects

The PRIR 2011 meeting is grateful for the support from the following sponsors and organisations:



### Effect of nitrogen fertilisation of strawberry plants on the efficacy of defence-stimulating biocontrol products against *Botrytis cinerea*

Presenting author: **Philippe C. Nicot** ([philippe.nicot@avignon.inra.fr](mailto:philippe.nicot@avignon.inra.fr))

Authors: Nicot PC<sup>1</sup>, Bardin M<sup>1</sup>, Lecompte F<sup>2</sup>

<sup>1</sup>INRA, UR407 Pathologie végétale, Domaine Saint Maurice, BP94, F-84143 Montfavet Cedex, France

<sup>2</sup>INRA, UR 1115 Plantes et Systèmes de Cultures Horticoles, Domaine Saint Paul, Site Agroparc, F-84914, Avignon Cedex

Nitrogen (N) fertilisation is known for its effect on the susceptibility of host plants to certain pests and pathogens and on the production of volatile secondary metabolites in response to attacks by herbivores. Although N is a key component in many compounds implicated in plant defence mechanisms, little is known on the possible effect of nitrogen fertilisation on the efficacy of defence-stimulating biocontrol agents. In the present work we examined the effect of five levels of N nutrition (0.5, 2, 5, 10 and 20 mMol L<sup>-1</sup> applied as nitrate in a hydroponic system) on the susceptibility of strawberry leaves to *Botrytis cinerea* and on the protective efficacy of Serenade Max (*Bacillus subtilis* QST713) and Chitoplant (chitosan), two biocontrol products presumed to induce plant defence mechanisms.

Two days after the application of the products, batches of leaf disks were excised, inoculated with *B. cinerea* and incubated in conditions conducive to disease development. The resulting lesions were photographed and their surface was assessed with the help of image analysis software. Plant fertilisation had a highly significant ( $p < 0.01$ ) effect on disease severity for both strains of *B. cinerea* tested, with lesion sizes smallest and largest, respectively, on leaves from plants with the lowest and highest N levels. Similar effects were observed with plants treated with either biocontrol product.

In addition, plant fertilisation significantly influenced the efficacy of the biocontrol products. Compared to the untreated control, Serenade Max provided significant protection against both strains of *B. cinerea* on plants with low levels of N fertilisation (0.5 and 2 mMol L<sup>-1</sup>), but not on those that received higher doses. In contrast, Chitoplant did not provide any significant protection against aggressive strain BC1. It provided a high level of protection (greater than 50%,  $p < 0.01$ ) against mildly aggressive strain BC21, but only for plants with low N fertilisation levels (0.5 and 2 mMol L<sup>-1</sup>).

Possible hypotheses and the relevance of these results for integrated protection will be discussed.