



Specific purity of hybrid larch Forest Reproductive Materials: how much does it matter?

Gwenaël Philippe, Stéphane Matz, Corinne Buret, Luc Pâques

► To cite this version:

Gwenaël Philippe, Stéphane Matz, Corinne Buret, Luc Pâques. Specific purity of hybrid larch Forest Reproductive Materials: how much does it matter?. Larix 2012;8. International Symposium of IUFRO Working Group S2.02.07 – Larch Breeding and Genetic Resources, International Union of Forest Research Organisations (IUFRO). AUT., Sep 2012, Hallormsstaðaskógur National Forest, Iceland. hal-02745016

HAL Id: hal-02745016

<https://hal.inrae.fr/hal-02745016>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



Specific purity of hybrid larch FRM: How much does it matter?

G. Philippe, S. Matz, C. Buret, L.E. Pâques

Pour mieux
affirmer
ses missions,
le Cemagref
devient Irstea



www.irstea.fr

LARIX 2012: Larch in a warm climate

8th international symposium of IUFRO Working Group S2.02.07 – Larch
Breeding and Genetic Resources

Hallormsstaðaskógur National Forest, Iceland, 11 - 13 September 2012


Dunkeld estate (Scotland): a place of pilgrimage




Different kinds of orchards

Nb of genotypes		Female species	Planting design	Pollination
♀	♂			
few → dozens	few → dozens	EL or JL or both	EL & JL intimately mixed or alternating rows	open
1	1 → dozens	EL or JL	separate orchards	SMP

Inter-specific crossings ...




... but also intra-specific crossings



Several maternal clones:

Crossings among clones + self-pollination

Case of s.o. with a single maternal clone



Self-pollination only ...
assuming the s.o. is well isolated from external pollen





The crops of hybridisation orchards composed of a mixture of seeds

- hybrid seeds
 - pure species
 - * crossings among different clones
 - * selfing
- (+ hybrids or pure species due to pollen contamination)



Hybrid rate varies according to genetic, climatic and anthropic factors

- the number of genitors of the mother species
- contribution of EL and JL clones to pollen cloud 
- flowering overlap between EL & JL
 - flowering ability
 - climate during flower initiation
 - flower induction treatments
- 
 - clonal phenology, « selfing ability »
 - Winter T°C
 - SMP



→ Hybrid purity will fluctuate from one orchard to another but also from year to year



Species purity can now be quantified in any s.o.

Distinguishing hybrids to the naked eye impossible (seeds) or long (seedlings)

Pâques et al. (2006)

1990's: isozymes but restrictions

Bergmann, Ruetz (1987), Häcker, Bergmann (1991), Ennos, Tang Quian (1994)

2000's: molecular markers based on cytoplasmic DNA

Acheré et al. (2004)

→ **Highly variable HL% in orchard crops (10-90%)**



Objectives of the study

1- Effect of HL purity on **stand productivity and quality**

- * To what extent are hybrids superior to pure species?
- * Is there a threshold below which HL% should not fall to ensure convenient stand production?
- * Is the importance of HL% the same for any kind of orchard?

2- Consequences of fluctuating FRM species purity in **seed orchard testing**



Material and Methods

Principle:

- Determination of taxonomic status of trees present in two s.o. testing trials
- Comparison of hybrid and pure species populations within each variety
- Estimation of thinning effect on HL%

M&M: studied varieties

Variety	orchard	Planting year	Country	Mother species	Number clones (EL/JL)
Halle	Halle	1959	B	JL & EL	<u>15</u> / <u>15</u>
FP 237	Grund	1978	DK	EL	<u>1</u> /17
FH 201	Les Barres	1976	F	EL	<u>1</u> /FS
Vaals	Vaals-01	1969	NL	EL	<u>1</u> /26
Esbeek	Esbeek-01	1971	NL	EL	<u>1</u> /4
Maglehem	51 Maglehem	1956	S	JL	8/ <u>1</u>



open pollination in all the orchards except FH 201 (SMP)

M&M: Variety testing trials

13

Planting in Spring 1995

RCB design:

4 blocks x 6 varieties

(64 individuals per plot)

Site 1

800 m elevation

deep soil

granite

8°C, 1100 mm

JL

Site 2


1000 m elevation

shallow soil

limestone

6°C, 1500 mm

EL





Site 2






M&M: Taxa identification

- Bud collection at age 10 or 14
- Taxa identification (INRA) using markers based on cytoplasmic DNA (LL and F-13)
- Were determined:
 - * Site 1: Halle, FH 201, Esbeek, Maglehem (2 blocks)
 - * Site 2: id + Vaals and FP 237 (2 blocks)

M&M: Performance assessment



Adaptation – Growth – Stem form



Growth: circumference + height at age 6

Stem straightness



Score 1




Score 5



Basal sweep (age 13 - site 2)


Results: Proportion of hybrids



Results: hybrids vs pure species - Adaptation traits


No difference between taxa for adaptation traits

But hybrid % higher in site 2 (15% mortality) than in site 1 (4% mortality)




Results: hybrids vs pure species - Mortality

	Survival
EL	38%
HL	86% **




Results: hybrids vs pure species - Growth



$$Volume = C^2 H^2 / 8\pi (H-1.30) \quad (Pardé 1961)$$

Results: hybrids vs pure species - Growth



Maternal species

blue: JL

red: EL



Results: hybrids vs pure species – Growth



Hybrids are always more vigorous than pure species but their superiority varies according to the variety and the site

Two explanatory factors :

→ the number of maternal clones

→ the adaptation of mother species to the planting site

Results: hybrids vs pure species – Further growth



ability to grow + competition



Results: hybrids versus pure species – Stem form

- H/circumference

Hybrids are less slender than pure species in most of the varieties

- Stem straightness and basal sweep

Hybrids are somewhat less straight but differences rarely significant




Results: effect of 1st thinning on species purity

Thinning at age 10 (site 1) and 14 (site 2)

	Density (number of trees per ha)		
	at planting	before thinning	after thinning
Site 1	1667	1610	940
Site 2	1667	1440	790

**systematic
thinning**




x	x	x	x	x	x	x	x
x	x	x	x	x	x	x	x
x	x	x	x	x	x	x	x
x	x	x	x	x	x	x	x
x	x	x	x	x	x	x	x
x	x	x	x	x	x	x	x
x	x	x	x	x	x	x	x
x	x	x	x	x	x	x	x


**selective
thinning**

irstea


Effect of 1st thinning on specific purity



Simulation of 2nd and 3rd thinning



Simulation of 2nd and 3rd thinnings





Consequences for the forest owner

They depend on **forest owner strategy** and the **variety**:

Stem form objective: hybrid richness does not matter much
but nonsense to target only form with hybrid varieties

Timber production

HL ~ pure species → moderate loss of productivity

HL >> pure species → potentially high risk

but limited consequences if hybrid% > 60-70%

Biomass production requires 100% hybrids



Conclusion

- 1- Some so-called HL varieties have a low hybrid rate
 - abstain from collecting the cones in high risk years
 - set a minimal threshold of purity for commercialisation

- 2- Hybrid superiority varies with the variety
 - number of maternal clones + adaptation of mother species

- 3- Hybrid rate increases after thinnings based on vigour
 - But $HL\% < 100\%$ if low initial species purity → loss of income
 - FRM species purity should be $> 60-70\%$ in high risk varieties



Conclusion

NO SEED LOT SHOULD BE COMMERCIALISED OR SHARED FOR TESTING WITHOUT MENTIONING HYBRID PURITY

■

A landscape photograph showing a grassy field with several young, thin trees in the foreground. In the background, there is a dense forest of taller trees under a cloudy sky.

Thank you for your attention !

Study funded by the French Ministry in charge of forests

ONF Vay P301