



**HAL**  
open science

# Influence of ascorbate-recycling, light and temperature during tomato fruit ripening on ascorbate pool and ascorbate-degradation

Vincent Truffault, H el ene Gautier, Rebecca Stevens

► **To cite this version:**

Vincent Truffault, H el ene Gautier, Rebecca Stevens. Influence of ascorbate-recycling, light and temperature during tomato fruit ripening on ascorbate pool and ascorbate-degradation. 11. International POG Conference Reactive Oxygen and Nitrogen Species in Plants, Sep 2013, Varsovie, Poland. hal-02745080

**HAL Id: hal-02745080**

**<https://hal.inrae.fr/hal-02745080>**

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destin ee au d ep ot et  a la diffusion de documents scientifiques de niveau recherche, publi es ou non,  emanant des  tablissements d'enseignement et de recherche fran ais ou  trangers, des laboratoires publics ou priv es.

Conference Paper: [Influence of ascorbate-recycling, light and temperature during tomato fruit ripening on ascorbate pool and ascorbate-degradation.](#)

[V Truffault](#) · [H Gautier](#) · R Stevens

[\[Show abstract\]](#)

BioTechnologia. Journal of Biotechnology Computational Biology and Bionanotechnology, Warsaw; 01/2013

Conference: BioTechnologia. Journal of Biotechnology Computational Biology and Bionanotechnology, At Warsaw, Volume: vol. 94(2) pp. 199 2013

## Abstract

Ascorbate is a powerful antioxidant in plants. Ascorbate concentration depends on its biosynthesis, recycling and degradation; these are under genetic control and closely related to environmental conditions. Ascorbate recycling, controlled by the enzymes monodehydroascorbate reductase (MDHAR) and dehydroascorbate reductase (DHAR), ensures a high-rate turnover of the ascorbate pool. Transgenic lines silenced for an MDHAR gene in a cherry tomato cultivar have been generated. The objective of this study was to determine whether limiting turnover of the ascorbate pool may affect levels of its reduced form and its degradation products. Degradation occurs via DHA to assayed candidate compounds: oxalate, tartrate and threonate. In order to investigate the impact of the reduction in MDHAR activity under different conditions of light and temperature on ascorbate concentration and ascorbate-degradation products, we used off-vine tomato fruits. Mature green fruits were harvested and placed in environmental chambers at 12EC, 23EC or 32EC, in darkness or light. Results are discussed and shed light on the stability of the ascorbate pool in fruits under different environmental conditions.