



HAL
open science

The efficacy of biocontrol against *Botrytis cinerea* on tomato depends on the strain of the pathogen

Marc Bardin, Claire Troulet, Philippe C. Nicot

► **To cite this version:**

Marc Bardin, Claire Troulet, Philippe C. Nicot. The efficacy of biocontrol against *Botrytis cinerea* on tomato depends on the strain of the pathogen. 10. International Congress of Plant Pathology (ICPP), Aug 2013, Pekin, China. , Acta Phytopathologica Sinica, 43 (Supplement), 620 p., 2013, ICPP 2013 Abstracts. hal-02745221

HAL Id: hal-02745221

<https://hal.inrae.fr/hal-02745221>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

P03.046 The efficacy of biocontrol against *Botrytis cinerea* on tomato depends on the strain of the pathogen

M. Bardin, C. Troulet and P.C. Nicot

INRA, UR407 Pathologie végétale, Domaine Saint Maurice, BP94, F-84143 Montfavet Cedex, France

Email: marc.bardin@avignon.inra.fr

The objective of the study was to estimate the risk of loss of biocontrol efficacy towards plant pathogens. To this end, the protective efficacy of both biocontrol agents *Microdochium dimerum* and *Bacillus subtilis* (Serenade Max®) was evaluated on tomato against 41 strains of *Botrytis cinerea* differing in their geographic origin, host of isolation and level of aggressiveness. The protective efficacy of *M. dimerum* was evaluated on whole tomato plants after co-inoculation of pruning wounds with *B. cinerea* and *M. dimerum* at the desired dose. Lesion expansion was measured on the stem from the 3rd to the 7th day after inoculation. A very high level of efficacy against all strains was obtained when the biocontrol agent was used at the recommended dose. At a 10-fold reduced application dose, a wide range of sensitivity was observed and a significant correlation was observed between the level of aggressiveness of a strain to tomato and its sensitivity to the biocontrol agent. For *B. subtilis*, tomato leaves were first treated with the biocontrol agent at the desired dose, two days before inoculation with a mycelial agar plug of *B. cinerea*. The resulting lesions were assessed from the 2nd to the 4th day after inoculation. The efficiency of this product was also significantly influenced by the strain of *B. cinerea* but no correlation was observed between the aggressiveness of *B. cinerea* and the protection provided by the biocontrol agent. This study reveals the importance of considering several strains of the pathogen when evaluating the efficacy of biocontrol agents, to obtain a good representation of the pathogen population.

2012中国国际影响力优秀学术期刊

中国科技核心期刊

《CAJ-CD规范》执行优秀期刊

ISSN 0412-0914
CN11-2184/S

植物病理学报

ACTA PHYTOPATHOLOGICA SINICA

ZHI WU BING LI XUE BAO

VOL.43 NO. **增刊** 2013

Supplement



ICPP 2013
August 25–30,
Beijing, China

Bio-security, Food Safety and Plant Pathology

ABSTRACTS

**10th International
Congress of Plant Pathology**

中国植物病理学会
Chinese Society for Plant Pathology