



HAL
open science

A rapid tools for physiological study of apricot fruit

Sylvie Bureau, Barbara Gouble, Jean Marc Audergon, Catherine M.G.C. Renard

► **To cite this version:**

Sylvie Bureau, Barbara Gouble, Jean Marc Audergon, Catherine M.G.C. Renard. A rapid tools for physiological study of apricot fruit. 15. International Symposium, Jun 2011, Yerevan, Armenia. hal-02745371

HAL Id: hal-02745371

<https://hal.inrae.fr/hal-02745371v1>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**JUNE
20-24**

**XV International
Symposium
on Apricot Breeding
and Culture**



armenia apricot



Com

A RAPID TOOLS FOR PHYSIOLOGICAL STUDY OF APRICOT FRUIT?

Sylvie BUREAU^{1,2}, Barbara GOUBLE^{1,2}, Jean-Marc AUDERGON³, Catherine M.G.C. RENARD^{1,2}

¹INRA, UMR408 Sécurité et Qualité des Produits d'Origine Végétale, F-84000 Avignon

²Université d'Avignon et des Pays de Vaucluse, UMR408 Sécurité et Qualité des Produits d'Origine Végétale, F-84000 Avignon

³Unité de Génétique et d'Amélioration des Fruits et Légumes, INRA, Domaine Saint Maurice, BP94, F-84143 Montfavet

Two wavelength regions in the infrared range have been tested in our laboratory for apricot quality analysis. The near-infrared (NIR, 800-2500 nm) range has been applied on intact fruits and the mid-infrared (MIR, 4000-650 cm^{-1}) one on fruit slurries. The type of samples, fruits or slurries, is dependent on the depth of light penetration. These techniques appear suitable for the characterisation of sugar and acid composition such as soluble solids content, titratable acidity, levels of sucrose, glucose, fructose, malic acid and citric acid.

Good results are surprisingly obtained in MIR spectroscopy for the prediction of ethylene production rate ($\ln(\text{nmol.kg}^{-1}.\text{h}^{-1})$) on a very large phenotypic variability of apricot, including good correlations for calibration and validation ($R^2 = 0.81$ and 0.83 respectively) and acceptable errors of validation and prediction (0.86 and $0.84 \text{ nmol.kg}^{-1}.\text{h}^{-1}$ respectively).

We are now interested in two objectives:

1) Do models obtained on a large phenotypic variability in MIR spectroscopy allow the prediction of ethylene levels on different cultivars, harvested over years and issued from different orchards?

2) Is it possible to use the non-destructive NIR spectroscopy for studying the change of apricot fruit during their maturation and post-harvest?

Compared to the method of reference, FTIR spectroscopy has been shown to be fast and easy applicable. This technique can be adapted to routine analysis in apricot industries.

Corresponding author: Sylvie BUREAU, Email: sylvie.bureau@avignon.inra.fr