



**HAL**  
open science

## Comparative effects of two water-accommodated fractions from light and heavy crude oils on early life stage of zebrafish *Danio rerio*

P. Perrichon, H el ene Budzinski, Karyn Le Menach, P. Bustamante, Xavier Cousin

### ► To cite this version:

P. Perrichon, H el ene Budzinski, Karyn Le Menach, P. Bustamante, Xavier Cousin. Comparative effects of two water-accommodated fractions from light and heavy crude oils on early life stage of zebrafish *Danio rerio*. 3. Young Environmental Scientists (YES) Meeting, Feb 2013, Cracovie, Poland. pp.82, 2013, Interdisciplinary discourse on current environmental challenges. hal-02745378

**HAL Id: hal-02745378**

**<https://hal.inrae.fr/hal-02745378>**

Submitted on 20 Nov 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destin ee au d ep ot et  a la diffusion de documents scientifiques de niveau recherche, publi es ou non,  emanant des  tablissements d'enseignement et de recherche fran ais ou  trangers, des laboratoires publics ou priv es.

## Posters

Session: Aquatic ecotoxicology

AT16

### Comparative effects of two water-accommodated fractions from light and heavy crude oils on early life stage of zebrafish *Danio rerio*.

P. Perrichon<sup>1</sup>, H. Budzinski<sup>2</sup>, K. Le Menach<sup>2</sup>, P. Bustamante<sup>3</sup> and X. Cousin<sup>1,4</sup>

<sup>1</sup>IFREMER, Place Gaby Coll, BP7, F17137 L' Houmeau, France

<sup>2</sup>Université de Bordeaux 1 (EPOC/LPTC, UMR CNRS 5805, France)

<sup>3</sup>Littoral Environnement et Sociétés (LIENSs), UMR 7266, 2 rue Olympe de Gouges, F-17000 La Rochelle, France

<sup>4</sup>INRA LPGP, Campus de Beaulieu – Bâtiment 16A, 35042 Rennes, France

Correspondence : prescilla.perrichon@ifremer.fr

Aquatic environment is constantly threaten by introduction of anthropogenic pollutants. Among them, polycyclic aromatic hydrocarbons (PAHs) are widespread in these areas and can represent a threat to biological functions of the fishes. PAHs are in fact complex mixtures of large number of compounds with variable properties (incl. physical-chemical and toxicity). These properties depend on their origin, pyrolytic or petrogenic. They however share a low solubility in water making direct exposure in water impossible. The use of fish early life stage (ELS) to assess toxicity of chemicals or media is considered as *in vitro* assay and therefore allows reducing live animals tests in accordance with the European legislation. The main objective of this study is to evaluate the suitability of water-accommodated fractions (WAF) and zebrafish ELS to assess oils toxicity. For this purpose, zebrafish ELS were exposed to two WAFs, one prepared with from light (Arabian Light) and heavy (Erika) crude oils to determine at several concentrations. Toxicity assessment was performed using lethal and sublethal (e.g. DNA damage, deformations, hatching rate, heart rate, swimming activity) endpoints recorded at different developmental stages. Combination of physiological, morphological and behavioral readouts indicated that WAF prepared from heavy oil is several orders of magnitude more toxic than light oil and confirmed the suitability of this approach.

**Keywords:** water-accommodated fraction – crude oils – bioassay – zebrafish – early life stage – embryo toxicity – endpoints



3RD YOUNG ENVIRONMENTAL  
SCIENTISTS MEETING KRAKÓW  
POLAND 11-13 FEBRUARY 2013



**“INTERDISCIPLINARY DISCOURSE ON  
CURRENT ENVIRONMENTAL CHALLENGES”**

**11 - 13 February 2013**



**Students Advisory Council**  
Speaking tube of SETAC Europe students