



HAL
open science

Monitoring viable airborne inoculum of *Botrytis cinerea* in the South-East of France over 3 years: relation with climatic parameters and the origin of air masses

Christel Leyronas, Philippe C. Nicot

► **To cite this version:**

Christel Leyronas, Philippe C. Nicot. Monitoring viable airborne inoculum of *Botrytis cinerea* in the South-East of France over 3 years: relation with climatic parameters and the origin of air masses. 10. International Congress of Plant Pathology (ICPP), Aug 2013, Pekin, China. 620 p. <hal-02745614>

HAL Id: hal-02745614

<https://hal.inrae.fr/hal-02745614v1>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



HAL Authorization

001.006 Monitoring viable airborne inoculum of *Botrytis cinerea* in the South-East of France over 3 years: relation with climatic parameters and the origin of air masses

C. Leyronas and P.C. Nicot

INRA, UR407 Pathologie végétale, Domaine Saint Maurice, CS 60094, F-84143 Montfavet cedex, France

Email: philippe.nicot@avignon.inra.fr

Viable airborne inoculum of *Botrytis cinerea* was monitored bimonthly during 3 years (September 2007- December 2010) on a site in the South-East of France located approximately 5 km away from susceptible crops. Viable inoculum was observed on 96 % of the sampling days, including during cold winter periods and hot and dry summer conditions. The concentration of airborne inoculum was significantly higher during day-time than at night. Peaks of concentration were recorded at different periods each year (September-October in 2008, May in 2010). The abundance of viable inoculum was positively correlated with average daily relative humidity and negatively correlated with air temperature and solar radiation. The analysis of backward trajectories suggested that air masses originating from the North or the South brought more viable inoculum than those from the West. This study showed that susceptible crops may be at danger from viable inoculum of *B. cinerea* during all seasons of the year, but that risk prediction models could be developed on the basis of climatic conditions and the origin of air masses.

2012中国国际影响力优秀学术期刊

中国科技核心期刊

《CAJ-CD规范》执行优秀期刊

ISSN 0412-0914
CN11-2184/S

植物病理学报

ACTA PHYTOPATHOLOGICA SINICA

ZHI WU BING LI XUE BAO

VOL.43 NO. **增刊** 2013

Supplement



ICPP 2013
August 25–30,
Beijing, China

Bio-security, Food Safety and Plant Pathology

ABSTRACTS

**10th International
Congress of Plant Pathology**

中国植物病理学会
Chinese Society for Plant Pathology