



Somatic embryo development in maritime pine (*Pinus pinaster*): a 2-DE proteomic analysis

Marie-Anne Lelu-Walter, Alexandre A. Morel, Caroline Teyssier, Martine Beaufour, Isabelle Reymond, Claire C. Le Mette, Martine Cadène, Jean-François Trontin, Philippe Label

► To cite this version:

Marie-Anne Lelu-Walter, Alexandre A. Morel, Caroline Teyssier, Martine Beaufour, Isabelle Reymond, et al.. Somatic embryo development in maritime pine (*Pinus pinaster*): a 2-DE proteomic analysis. 4. UPSC-INRA (UPRA) Bilateral Meeting, Mar 2012, Umea, Sweden. 2012, 4th UPSC-INRA (UPRA) bilateral meeting. Meeting book with abstracts. hal-02746152

HAL Id: hal-02746152

<https://hal.inrae.fr/hal-02746152>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Somatic embryo development in maritime pine (*Pinus pinaster*):
a 2-DE proteomic analysis**

Alexandre Morel^{a*}, Caroline Teyssier^a, Martine Beaufour^c, Isabelle Reymond^b, Claire Le
Metté^a, Martine Cadene^c, Jean-François Trontin^b, Philippe Label^a, Marie-Anne Lelu-Walter^a

^a : INRA, UR 588 Research Unit: Breeding, Genetic and Physiology of Forest Trees

^b : FCBA Technical Institute, Biotechnology Lab, France

^c : CNRS UPR4301, Centre de Biophysique Moléculaire, France

*alexandre.morel@orleans.inra.fr

Summary (300 words)

Development of clonal propagation method, such as somatic embryogenesis (SE) has potentially numerous applications such as the production of a large number of genetically improved plants. However it is necessary to optimise somatic embryo development, which remains difficult in pine species. Since the early protocols of SE were developed for spruce species (*Picea*), it was assumed that they would be applicable to pines; however, it became apparent that pines were less responsive. One limitation to large-scale propagation of majority of commercial pine species through SE is the lack or low somatic embryo maturation efficiency, which result in the inability to capture certain families. This has caused great concerns among breeders and foresters of potential adverse selection pressure.

Current maturation protocols lead to a development of mature somatic embryos that morphologically resemble zygotic embryos. These somatic embryos are harvested after arbitrarily chosen periods of time and are germinated and further grown in a greenhouse. This procedure only gives an indication that the maturation was successful if produced somatic embryos were able to convert to plants. Such an empirical approach does not give any information of the quality of somatic embryos with respect to storage reserves accumulation nor does it gives information on the optimal time for harvesting to achieve maximal plant conversion rates. Therefore, there is a need to develop markers that could be used for quality control of different batches of somatic embryos that are matured, or when different maturation protocols are applied. Storage protein accumulation has been followed and the identity of the proteins accumulated by the somatic embryos has been tested by 2D gel electrophoresis.

Final objective is to have a better understanding of the maturation of *Pinus pinaster*. The description of SE would contribute to optimise the maturation process and the *in vitro* production of plants.