

Identification of QTL of transcripts (eQTL) and of proteins (pQTL) on foie gras of mule ducks

Christel Marie Etancelin, Yoannah François, Caroline Molette, Alain Vignal, Stephane Davail

▶ To cite this version:

Christel Marie Etancelin, Yoannah François, Caroline Molette, Alain Vignal, Stephane Davail. Identification of QTL of transcripts (eQTL) and of proteins (pQTL) on foie gras of mule ducks. 8. European Symposium on Poultry Genetic, Sep 2013, Venise, Italy. 2013. hal-02746170

HAL Id: hal-02746170 https://hal.inrae.fr/hal-02746170

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Identification of QTL of transcripts (eQTL) and of proteins (pQTL) on foie gras of mule

Y. François*1,2,3,4, C. Molette^{2,4}, C. Marie-Etancelin³, A. Vignal⁴, S. Davail¹

¹IPREM-EEM, IUT des Pays de l'Adour, BP 201-40004 Mont de Marsan, FRANCE ²INRA, INPT-ENSAT, INPT-ENVT, UMR TANDEM, 31326 Castanet-Tolosan, France ³INRA, UR 631. Station d'Amélioration Génétique des Animaux, 31326 Castanet-Tolosan, France ⁴INRA, INPT-ENVT, UMR444 Laboratoire de Génétique Cellulaire, 31326 Castanet-Tolosan, France


In France, 97% of the foie gras production comes from mule ducks, an inter-specific hybrid obtained by crossing a female common duck (*Anas Platyrhyncos*) and a Muscovy drake (*Cairina Moschata*). As mule ducks are infertile, the genetic improvement of the fatty liver quality has to be done through the genetic selection of parental lines. Thus, knowing the impact of genes on the quality traits could be useful for selection purposes. The project is based on a QTL (Quantitative Trait Loci) experimental design with more than 1,500 phenotyped mule ducks. In 2013, Kileh-Wais *et al.* found 11 significant QTL related to liver quality traits, with QTL clusters on chromosomes 2 and 9. Focusing on these QTL, proteomic and transcriptomic analyzes were performed on fatty livers of a subset of 300 mule ducks. Proteins were quantified using the bi-dimensional gel electrophoresis method and transcripts were quantified by quantitative PCR (Fluidigm® technology). The regulatory networks implicated in the fatty liver quality are presently under investigation at both levels through the identification of pQTL and of eQTL, ie QTL controlling protein and transcript levels, respectively.

Proceedings of the

8th European Symposium on Poultry Genetics

World's Poultry Science Association (WPSA)

Working Group 3 "Breeding and Genetics"



25-27 September 2013

San Servolo Conference Centre

Venice, ITALY

www.epgs2013.com