



# VPgs (Viral Proteins Genome-Linked) from phytoviruses display intrinsically disordered features

Jocelyne J. Walter, Eugénie Hébrard, Thierry Michon

## ► To cite this version:

Jocelyne J. Walter, Eugénie Hébrard, Thierry Michon. VPgs (Viral Proteins Genome-Linked) from phytoviruses display intrinsically disordered features. 13. Rencontres de virologie végétale, Jan 2011, Aussois, France. hal-02746310

**HAL Id: hal-02746310**

**<https://hal.inrae.fr/hal-02746310>**


Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# 13<sup>èmes</sup> rencontres de virologie végétale



Aussois du 16 au 20 janvier 2011



## 7- VPgs (Viral Proteins Genome-Linked) from phytoviruses display intrinsically disordered features

Jocelyne WALTER<sup>1</sup>, Eugénie HEBRARD<sup>2</sup>, Thierry MICHON<sup>1</sup>

<sup>1</sup> UMR GDPP, Institut de Biologie Végétale Moléculaire, Villenave d'Ornon, France

<sup>2</sup> UMR 1097, Institut de Recherche pour le Développement (IRD), BP 64501, 34394 Montpellier cedex 5, France,

VPgs are viral proteins linked at the 5' end of the genome of many plant and animal viruses. Although sharing this common feature, VPgs are highly diverse in size, sequence and function. They are involved in key steps of the viral cycle such as genome replication, protein translation and viral movement in plant hosts. According to *in silico* predictions, many VPgs of non-related viruses share intrinsic disorder properties. These predictions were experimentally validated for several potyviral and sobemoviral VPgs. In this study, we show that the monomeric LMV and RYMV VPgs display structural features inducing an hydrodynamic behavior in-between a native premolten globule and a molten globule. Secondary structure elements are predicted *in silico* and detected by Far UV-circular dichroism. Transitions from disorder to order were observed using low percentages of TFE.

Intrinsically disordered domains of phytovirus VPgs generally comprised the functional sites involved in plant resistance-breaking. For instance, the central domain of the LMV VPg predicted as intrinsically disordered is involved in VPg association with the natively folded plant resistance factor eIF4E. One may expect that during the complex formation VPg undergoes a disorder-to-order transition. Fast kinetic measurements of the association between eIF4E and the VPg central domain are consistent with a two steps association compatible with the "fly-casting" model.