HAL open science

Heavy metal availability and their relationships with soil microbial characteristics in agricultural carbonated soils

Ana de Santiago Martin, Nathalie Cheviron, Jose R. Quintana, Concepción González Huecas, Antonio López Lafuente, Christian Mougin

To cite this version:

Ana de Santiago Martin, Nathalie Cheviron, Jose R. Quintana, Concepción González Huecas, Antonio López Lafuente, et al.. Heavy metal availability and their relationships with soil microbial characteristics in agricultural carbonated soils. 6th International Symposium of Interactions of Soil Minerals with Organic Components and Microorganisms (ISMOM) Soil Interfaces in a Changing World, Jun 2011, Montpellier, France. 2011. hal-02746600

HAL Id: hal-02746600

https://hal.inrae.fr/hal-02746600v1

Submitted on 15 Mar 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

HEAVY METAL AVAILABILITY AND THEIR RELATIONSHIPS WITH SOIL MICROBIAL CHARACTERISTICS IN AGRICULTURAL CARBONATED SOILS

Ana de Santiago [a], Nathalie Cheviron b, José R. Quintana [a], Concepción González [a], Antonio L. Lafuente [a], Christian Mougin [b]
[a] Departamento de Edafología - Facultad de Farmacia - Universidad Complutense de Madrid (UCM) - Plaza de Ramón y Cajal s/n - 28040 Madrid - Spain
[b] UPR 251 - PESSAC - Physico-chimie et Ecotoxicologie des sols d'Agrosystèmes Contaminés - INRA - Route de St-Cyr - 78026 Versailles - France

INTRODUCTION

Soil contamination by heavy metals (HMs) can have negative long-term toxic effects on ecosystems affecting key microbial processes and decreasing the number and activity of soil microorganisms.
Soil enzyme activities are recognized as a highly sensitive bio-indicator to HMs and the methodology for their determination is rapid, simple and inexpensive. However, natural variability within and between soils could be a decisive impediment to use simple parameters such as single enzyme activities, suggesting the use of the combination with other biochemical properties such as organic C, total N, soil microbial biomass or total DNA load.

The aim of this study was to: i) assess the impact of heavy metals on soil enzyme activities (arylsulfatase, alkaline phosphatase, ß-galactosidase, urease and dehydrogenase) and soil DNA loads (total, bacterial and fungal DNA) and ii) evaluate the microbiological parameters measured as HMs bio-indicators and their relationships with several HM availability indexes.



MATERIAL AND METHODS

Soil sampling and contaminant experiment

• 10 unpolluted agricultural carbonated soils were collected down to 30 cm depth in Madrid (Spain).
• Soils were spiked with Cd, Cu, Zn and Pb at two contamination levels, corresponding to current EU legislative limits for HMs in agricultural soils -T1- and semi-maximum concentrations for sewage sludge -T2- (Table), and after soils were subjected to wetting-drying cycles for 365 contact days. [Soil samples without addition of HMs were employed as a control = T0]

mg Kg^{-1}	Cd	Cu	Zn	Pb
T1	3	140	300	300
T2	20	875	2000	600

Physicochemical analyses

• Soil properties related with availability of heavy metals were measured such as pH, particle size, CEC, equivalent $CaCO_3$, Fe-Mn oxides, available P, total N and several parameters related with the organic matter.

• Availability indexes: 0.1 M $NaNO_3$, 10 mM LMWOA and 0.005 M DTPA.

Microbiological analyses

• Soil enzyme activities: dehydrogenase (DEH), arylsulfatase (ARYS), β-galactosidase (BGAL), urease (URE) and alkaline phosphatase (ALP) activities using standardized methods developed by the Unité PESSAC (INRA Versailles-Grignon).

• Soil DNA loads:
- DNA extraction: by Fast DNA kit (MP Biomedical).
- DNA quantification:
  - Total DNA load (tDNA): absorbance at 260 nm.
  - Fungal and bacterial DNA loads (fDNA and bDNA): qPCR.

RESULTS AND CONCLUSIONS

Effects of HMs on microbiological parameters



BGAL and URE activities decreased significantly in both contamination levels. ARYS, ALP and DEH were significantly inhibited T2 level.
Total DNA load was significantly lower in T1 and T2 than in T0. Fungal DNA load showed no significant differences between treatments, while bacterial DNA load decreased significantly in both T1 and T2.

Relationship between microbial parameters, availability indexes and soil properties (RDA analysis)



Microbial parameters distribution pattern was mainly related to DTPA Cd, Cu and Zn and LMWOA Pb.

Fe oxides and recalcitrant organic C (ROC) were the main soil properties that determined samples dispersion.

Relationship between some selected microbiological ratios and availability indexes

The Pearson's correlation analysis showed no clear effects of contaminants on DNA loads and enzyme activities. In contrast, simple biological ratios were significant correlated with extractable metals.



ACKOWLEDGEMENTS: *This work was supported by the **Red CARESOIL** (Ref. P2009/AMB-1648) and the **Unité PESSAC** (INRA Versailles-Grignon)*