

## Apricot core collection assessment using morphological characterization

Lamia Krichen, Hedia Bourguiba, Jean Marc Audergon, Neïla Trifi-Farah

► **To cite this version:**

Lamia Krichen, Hedia Bourguiba, Jean Marc Audergon, Neïla Trifi-Farah. Apricot core collection assessment using morphological characterization. 15. International Symposium, Jun 2011, Yerevan, Armenia. hal-02747317

**HAL Id: hal-02747317**

**<https://hal.inrae.fr/hal-02747317>**

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.




**JUNE  
20-24**

**XV International  
Symposium  
on Apricot Breeding  
and Culture**



*armenia apricot*





## APRICOT CORE COLLECTION ASSESSMENT USING MORPHOLOGICAL CHARACTERIZATION

**Krichen Lamia<sup>1</sup> Bourguiba Hedia<sup>1</sup>, Audergon Jean-Marc<sup>2</sup>, Trifi-Farah Neila<sup>1</sup>**

<sup>1</sup> Laboratoire de Génétique Moléculaire, Immunologie et Biotechnologie. Faculté des Sciences de Tunis, Campus Universitaire – Tunis-El Manar, 2092, Tunisie.

<sup>2</sup> INRA UR1052. Unité de Génétique et Amélioration des Fruits et Légumes, Centre de Recherches d'Avignon - Domaine St Maurice –BP94, 84143 Montfavet Cedex, France.

To study the morphological variability of Tunisian apricot germplasm, 112 accessions corresponding to seventy six grafted cultivars and thirty six Bargougs propagated by seeds were collected from the principal areas of apricot cultivation. Thirty-four morphological characters concerning trees, leaves and fruits, were exploited to analyze the phenotypic variability. All the studied characters were polymorphic and showed a large variability among the Tunisian apricot germplasm. Using maximization strategy algorithm (Gouesnard et al. 2001), data analysis permitted the construction of a core collection allowing the selection of a twenty three accessions representative of the Tunisian genetic variability. This core collection expressed 100% of the global diversity for twenty six characters, more than 92% of the global diversity for six quantitative characters and 75% for two characters. Such results demonstrated that our core collection is well representative of the studied germplasm.

Corresponding author: Lamia KRICHEN, email: [krichlam@planet.tn](mailto:krichlam@planet.tn)