



HAL
open science

Heterogeneity of subspecies *Mycobacterium avium* paratuberculosis from genotype to phenotype

Louise Lefrançois, Christelle Bodier, Thierry Cochard, Florence Gilbert, Sylvie Canepa, S Lecher, D Raze, Philippe Lanotte, E. Haguenoer, I.A. Sevilla, et al.

► **To cite this version:**

Louise Lefrançois, Christelle Bodier, Thierry Cochard, Florence Gilbert, Sylvie Canepa, et al.. Heterogeneity of subspecies *Mycobacterium avium* paratuberculosis from genotype to phenotype. 11. International Colloquium on Paratuberculosis, Feb 2012, Sydney, Australia. pp.245, 2012. hal-02748073

HAL Id: hal-02748073

<https://hal.inrae.fr/hal-02748073>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

HETEROGENEITY OF SUBSPECIES *MYCOBACTERIUM AVIUM PARATUBERCULOSIS* FROM GENOTYPE TO PHENOTYPE

Lefrancois LH¹, Bodier CC¹, Cochard T¹, Gilbert F¹, Canepa S², Lecher S³, Raze D³, Lanotte P⁴, Haguenoer E⁴, Sevilla IA⁵, Stevenson K⁶, Behr M⁷, Collins D⁸, Loch C³, Biet F

1 INRA, ISP Centre de Tours, Nouzilly, France

2 INRA, PAIB, UMR6175 Centre de Tours, Nouzilly, France

3 CIIL, INSERM, CNRS, University of Lille Nord, Institut Pasteur, Lille, France

4 CHRU Bretonneau, Tours, France


5 Neiker-tecnalia, Bizkaia, Spain

6 Moredun Research Institute, Penicuik, Scotland, United Kingdom

7 McGill University, QC, H3G 1A4 Montreal, Canada

8 AgResearch, National Centre for Biosecurity and Infectious Disease, Wallaceville, New Zealand

Background – In the subspecies *M. avium* ssp. *paratuberculosis* (Map) two groups, known as Cattle (C) and Sheep (S), have been defined by genotyping. Recent studies show that Map C and S have different phenotypes with respect to infection of macrophages and iron metabolism. Map is adapted to the gastrointestinal tract of ruminant, but the mechanism of entry is currently unknown. In this study, we investigated the phenotype of the Map-host interaction, involving the virulence factor heparin-binding hemagglutinin (HBHA), for both groups of Map. HBHA is described in *M. tuberculosis* as a major adhesin required for extrapulmonary dissemination of the tubercle bacillus. Method – A large collection of Map isolates (types C & S) were genotyped by MIRU-VNTR and RFLP-IS900. The polymorphism of the hbha gene was investigated by fragment analysis using GeneMapper technology. Structure-functions properties of recombinant HBHA (types C & S) were analyzed by Heparin- Sepharose chromatography and SPR analysis based on Biacore technology. Results – In silico analyses of both types of Map have revealed two forms of hbha. This observation, showing that hbha is distinct according to the group, was confirmed using GeneMapper on 83 Map strains (65 Map C & 18 Map S) with various genotypes. We found that Map type C produces HBHA with a short C-terminal domain, while that of type S presents a long C-terminal domain, similar to that of HBHA produced by *M. tuberculosis*. The purification of HBHA from Map type C and S by Heparin-sepharose chromatography highlighted a correlation between their affinities to heparin and the length of their C-terminal domain confirmed by Biacore analysis. Conclusion – We show for the first time that the types C and S of Map may be distinguished by the type of HBHA they produce, which differs in size and adherence properties. Thus, HBHA participates in the genotypic and phenotypic differences observed between the C and S types of Map.



11th International
Colloquium on
Paratuberculosis

Proceedings of the 11th International Colloquium on Paratuberculosis



THE UNIVERSITY OF
SYDNEY

SYDNEY, AUSTRALIA
5 – 10 FEBRUARY 2012



**INTERNATIONAL ASSOCIATION
FOR PARATUBERCULOSIS**