



Metabolite profiling and feeding bioassays suggest a major role for a dicaffeoylquinic acid in induced resistance of peach to *Myzus persicae* aphid

Jean-Luc Poessel, Marie-Hélène Sauge, Michael Staudt, Claire Dufour, Catherine Deborde, Yvan Rahbé, B. Jackson, Cédric Renaud, Mickael M. Maucourt, Marie Noëlle Corre, et al.

► To cite this version:

Jean-Luc Poessel, Marie-Hélène Sauge, Michael Staudt, Claire Dufour, Catherine Deborde, et al.. Metabolite profiling and feeding bioassays suggest a major role for a dicaffeoylquinic acid in induced resistance of peach to *Myzus persicae* aphid. PRIR 2011 Joint meeting of the "PR-proteins Workshop" and the "Working Group Induced Resistance in Plants Against Insects and Diseases", Sep 2011, Neuchatel, Switzerland. 154 p., 2011, PR-Proteins and Induced Resistance Against Pathogens and Insects. Abstracts book. hal-02748156

HAL Id: hal-02748156

<https://hal.inrae.fr/hal-02748156>

Submitted on 3 Jun 2020



HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



PR-Proteins and Induced Resistance Against Pathogens and Insects

molecular biology meets application

 Neuchâtel, Switzerland
 4-8 September 2011

Abstract|book



Metabolite profiling and feeding bioassays suggest a major role for a dicaffeoylquinic acid in induced resistance of peach to *Myzus persicae* aphid

Presenting author: **Jean-Luc Poëssel** (Jean-Luc.Poessel@avignon.inra.fr)

Authors: Poëssel JL¹, Sauge MH², Staudt M³, Dufour C⁴, Deborde C^{5,6}, Rahbé Y⁷, Jackson B³, Renaud C⁸, Maucourt M^{6,9}, Corre MN¹, El-Aouni H³, Lacroze JP², Moing A^{5,6}

¹INRA, UR 1052, Génétique et Amélioration des Fruits et Légumes, Avignon, France

²INRA, UR 1115, Plantes et Systèmes de culture Horticoles, Avignon, France

³CEFE-CNRS, Montpellier, France

⁴INRA, UMR 408, Sécurité et Qualité des Produits d'Origine Végétale, Avignon, France

⁵INRA, UMR 1332 Biologie du Fruit et Pathologie, Bordeaux, France

⁶Plateforme Métabolome du CGFB, IBVM, Bordeaux, France

⁷INRA, UMR 203, Biologie Fonctionnelle, Insectes & Interactions, Lyon, France

⁸INRA, UR 419, Unité de Recherches sur les Espèces Fruitières, Bordeaux, France

⁹Université de Bordeaux, UMR 1332 Biologie du Fruit et Pathologie, France

The green peach aphid (GPA, *Myzus persicae* Sulzer) is a generalist insect pest infesting agricultural and horticultural crops worldwide. GPA causes significant direct damages to peach, its primary host, and may transmit viruses that cause serious diseases such as Sharka conferred by the *Plum Pox Potyvirus*. Resistant genotypes have been identified among the peach germplasm and are used in breeding programs to create peach cultivars showing a durable resistance to GPA. Among these genetic resources, Rubira, a red-leaf cultivar used as rootstock and carrying the dominant resistance gene *Rm2*, shows a strong and induced antixenosis-type resistance associated with hypersensitive-like necrotic lesions on developing leaves.

To study the mechanisms involved in induced resistance of Rubira to GPA, we used a metabolomics approach and investigated metabolite changes occurring in shoot apices of Rubira and a susceptible cultivar after a 48-hours infestation with GPA. Untargeted ¹H NMR and targeted LC (sugars, organic acids, amino acids), LC-MS (secondary metabolites including phenolic and cyanogenic compounds) and GC-MS (Volatile Organic Compounds, VOCs) were applied. While no significant modifications occurred in the susceptible cultivar, dramatic changes in primary and secondary metabolites were observed in Rubira following infestation. Carbohydrates and most organic acids showed a marked decrease. Several amino acids, including lysine and branched-chain and aromatic amino acids, showed a large accumulation whereas levels of glutamine, proline and threonine were greatly reduced. Infestation of Rubira by GPA also triggered the release of VOCs, mainly methyl-salicylate and (E,E)- α -farnesene, and the accumulation of secondary metabolites. Chlorogenic acid (5-caffeoylquinic acid, 5-CQ) and 3,5-dicaffeoylquinic acid (3,5-diCQ) were the main phenolics of shoot apices. Strikingly, these closely related compounds showed differential responses to infestation: while 5-CQ level did not change, 3,5-diCQ showed a significant accumulation and therefore could be involved in resistance. The effect of 5-CQ and 3,5-diCQ on GPA was further studied in bioassays with artificial diets. Whereas 5-CQ did not show any effect on larval development, 3,5-diCQ was highly toxic and larvae did not survive at the concentration as low as 1mM (516 mg.L⁻¹). These results suggest that 3,5-diCQ could play a major role in antixenosis resistance of Rubira to green peach aphid.