



**HAL**  
open science

## Use of volatile compound metabolic signatures in poultry tissues to back-trace dietary exposure to xenobiotics

Erwan Engel, Jérémy Ratel, Philippe Berge, Weeraya Khummueng, Cyril Feidt, Brigitte Roudaut, Bruno Le Bizec

### ► To cite this version:

Erwan Engel, Jérémy Ratel, Philippe Berge, Weeraya Khummueng, Cyril Feidt, et al.. Use of volatile compound metabolic signatures in poultry tissues to back-trace dietary exposure to xenobiotics. 3. MoniQA International Conference, Sep 2011, Varna, Bulgaria. 1 p., 2011, 3èmes MoniQA International Conference. hal-02748205

**HAL Id: hal-02748205**

**<https://hal.inrae.fr/hal-02748205v1>**

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Use of Volatile Compound Metabolic Signatures in Poultry Tissues to Back-Trace Dietary Exposure to Xenobiotics

E. Engel<sup>1,\*</sup>, J. Ratel<sup>1</sup>, P. Berge<sup>1</sup>, W. Khummueng<sup>1,2</sup>, C. Feidt<sup>3</sup>, B. Roudaut<sup>4</sup>, B. Le Bizec,<sup>5</sup>

<sup>1</sup> INRA, UR370 QuaPA, Saint-Genès-Champanelle, France.

<sup>2</sup> Science Department, Prince of Songkla University, Pattani campus, Thailand

<sup>3</sup> UR AFPA, USC340 INRA - Nancy Université, France.

<sup>4</sup> ANSES, Fougères, France

<sup>5</sup> ONIRIS, Nantes, France

**ABSTRACT** We investigated the feasibility of using volatile compound signatures of liver in poultry to detect previous dietary exposure to different types of xenobiotic. Six groups of broiler chickens were fed a similar diet either non-contaminated or contaminated with polychlorinated dibenzo-*p*-dioxins/-furans (PCDD/Fs), polychlorinated biphenyls (PCBs), polybrominated diphenyl ethers (PBDEs), polycyclic aromatic hydrocarbons (PAHs) or coccidiostats. The liver of each chicken was analysed by solid-phase microextraction - mass spectrometry (SPME-MS) for volatile compound metabolic signature and by gas chromatography - high resolution mass spectrometry (GC-HRMS), gas chromatography - tandem mass spectrometry (GC-MS/MS) and liquid chromatography - tandem mass spectrometry (LC-MS/MS) to quantify xenobiotic residues. The results show that the volatile compound metabolic signature could clearly differentiate the non-contaminated chickens from those contaminated with PBDEs, PAHs or coccidiostats. The results for PAHs showed a clear metabolic response in the liver although these rapidly metabolized xenobiotics are undetectable in this organ by the targeted reference analytical method. However, the rough metabolic signature obtained by SPME-MS did not enable us to evidence previous exposure to slowly metabolized compounds such as PCDD/Fs and PCBs, the residues of which are clearly detected by targeted reference methods.

**KEYWORDS** : Non targeted approach, environmental micropollutant, drug, liver, poultry