



HAL
open science

Dynamics of DNA methylation levels in maternal and paternal rabbit genomes after fertilization

Adriana Reis E Silva, Juliette Salvaing, Pierre Adenot, Nathalie N. Daniel, Catherine Archilla, Nathalie N. Peynot, C.M. Lucci, Nathalie Beaujean, Véronique Duranthon

► To cite this version:

Adriana Reis E Silva, Juliette Salvaing, Pierre Adenot, Nathalie N. Daniel, Catherine Archilla, et al.. Dynamics of DNA methylation levels in maternal and paternal rabbit genomes after fertilization. 22. Wilhelm Bernhard Workshop, International Federation for Cell Biology. USA., Aug 2011, Riga, Latvia. hal-02748414

HAL Id: hal-02748414

<https://hal.inrae.fr/hal-02748414>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

WILHELM BERNHARD

WORKSHOP RIGA 2011

XII

Program & Abstracts

22nd Wilhelm Bernhard Workshop

August 25 - 29, 2011

Riga - LATVIA

Chromatin - Nuclear matrix - Nucleolus
Transcription - DNA and Drug Interactions
Apoptosis - DNA Repair - Epigenetics

INFORMATION 22wbw@lu.lv



ERAF
EIROPAS REĢIONĀLĀS
ATĪSTĪBAS FONDS

DYNAMICS OF DNA METHYLATION LEVELS IN MATERNAL AND PATERNAL RABBIT GENOMES AFTER FERTILIZATION

A.R. Reis e Silva^{1,2}, J.Salvaing¹, P.Adenot¹, N.Daniel¹, C. Archilla¹, N.Peynot¹, C.M.Lucci², N.Beaujean¹, V.Duranthon¹

1. INRA, UMR 1198 Biologie du Développement et Reproduction, F-78350 Jouy en Josas, France;

2. Faculty of Veterinary Medicine, University of Brasilia, Brasilia, DF 70910-900, Brazil; E-mail: Juliette.Salvaing@jouy.inra.fr

The reprogramming of DNA methylation in early embryos has been considered to be essential for the reprogramming to totipotency, the embryonic genome activation (EGA) and subsequent development. However, its degree appears to differ as a function of species and it may be altered by the in vitro environment. While the rabbit is a pertinent model for species with a delayed EGA because both in vivo and in vitro developed embryos are easily available, the status of DNA methylation levels in both parental genomes after fertilization remains controversial. In order to generate precise data on the DNA methylation status in rabbit zygotes, we first of all defined five pronuclear (PN) stages during the first cell cycle, which we subsequently used to classify in vivo and in vitro developed rabbit zygotes. This allowed us to precisely quantify the methylated DNA during the one cell stage, as well as the total DNA content, which was used for normalization. We thus showed that both pronuclei display distinct patterns of DNA methylation reprogramming. In the maternal pronucleus (MP) the methylation level remains constant throughout the one cell stage, thanks to maintenance methylation during the S-phase. Conversely, in the paternal pronucleus (PP) partial demethylation occurs before replication, probably as a result of active DNA demethylation, while maintenance methylation subsequently takes place during the S-phase. Interestingly, we showed that PP DNA methylation reprogramming is partially altered by the in vitro environment. Taken together, our approach evidences that rabbit is one of the species displaying partial DNA demethylation in the PP, and for the first time demonstrates maintenance methylation activity in both pronuclei during the first S phase.

022