



## Bayesian modeling of the dynamics of complex of salmon populations in the North Atlantic

Félix Massiot-Granier, Etienne Prévost, Gerald Chaput, Ted Potter, Gordon Smith, Atso Romakkaniemi, Etienne Rivot

### ► To cite this version:

Félix Massiot-Granier, Etienne Prévost, Gerald Chaput, Ted Potter, Gordon Smith, et al.. Bayesian modeling of the dynamics of complex of salmon populations in the North Atlantic. International Statistical Ecology Conference, Jul 2012, Krokkeiva, Norway. hal-02748883

**HAL Id: hal-02748883**

**<https://hal.inrae.fr/hal-02748883>**

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## **Bayesian modeling of the dynamics of complex of salmon populations in the North Atlantic**

Félix Massiot-Granier<sup>1,2</sup>, Prevost Etienne<sup>2</sup>, Chaput Gerald<sup>3</sup>, Potter Ted<sup>4</sup>, Smith Gordon<sup>5</sup>, Romakkaniemi Atso<sup>6</sup>, Rivot Etienne<sup>1</sup>

<sup>1</sup>*Agrocampus-Ouest, INRA, UMR ESE, Rennes, France*, <sup>2</sup>*INRA, UMR Ecobiop, St Pee sur Nivelle, France*, <sup>3</sup>*Fisheries & Oceans Canada, Moncton, Canada*, <sup>4</sup>*CEFAS, fisheries lab, Lowestoft, Suffolk, UK*, <sup>5</sup>*FRS FL field station, Montrose, Scotland, UK*, <sup>6</sup>*Finnish Game & Fisheries Res Inst, Oulu, Finland*

Life cycle models are appropriate tools to evaluate multiple factors affecting wild animal populations at different life stages and at various spatial and temporal scales. They allow us to unravel the underlying mechanisms and predict the response of populations under plausible future scenarios.

Bayesian state-space modelling is appropriate for embedding complex demographic models within a statistical framework accommodating various sources of data, to derive inferences together with a fair appraisal of the uncertainty around estimates and predictions.

This multiscale approach was used to model Atlantic salmon population dynamics in the northern Atlantic. The model captures the complex meta-population structure stemming from homing behaviour for reproduction in freshwater. The different population components share some common environmental conditions and harvest pressure during their long migration in the sea.

The model was fitted to a 40 years data base of catches and homewater returns available for each country contributing to the population complex. The multiscale approach allows us to explore long term trends and climate influence on key population parameters shared by several population components, such as marine survival, together with time and spatial variability of region-specific life history traits characterizing the freshwater phase of the life cycle.