



HAL
open science

Surprising effect of nitrogen nutrition regimes on the development of *Botrytis cinerea* and *Sclerotinia sclerotiorum* on lettuce plants

Manzoor Ali Abro, François Lecompte, Magali Duffaud, Philippe C. Nicot

► To cite this version:

Manzoor Ali Abro, François Lecompte, Magali Duffaud, Philippe C. Nicot. Surprising effect of nitrogen nutrition regimes on the development of *Botrytis cinerea* and *Sclerotinia sclerotiorum* on lettuce plants. 9. Rencontres de Phytopathologie-Mycologie de la Société Française de Phytopathologie, Jan 2012, Aussois, France. 2012, Journées Jean Chevaugéon 2012. hal-02749125

HAL Id: hal-02749125

<https://hal.inrae.fr/hal-02749125>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Surprising effect of nitrogen nutrition regimes on the development of *Botrytis cinerea* and *Sclerotinia sclerotiorum* on lettuce plants

M.A. Abro (1), F. Lecompte (2), M. Duffaud (1) and P.C. Nicot (1)

(1) INRA, UR407 Pathologie végétale, Domaine Saint Maurice, BP94, F84143 Montfavet cedex, France.

(2) INRA, UR 1115 Plantes et Systems de Cultures horticoles, Domaine Saint Paul, Site Agroparc, F-84194, Avignon cedex, France

Although nitrogen (N) fertilisation is known to affect disease development in a variety of host-parasite systems, information is lacking on lettuce drop and grey mould, caused respectively by *Sclerotinia sclerotiorum* and *Botrytis cinerea*. Studies were carried out to investigate the effect of five regimes of nitrogen nutrition on the susceptibility of lettuce plants to six strains of *B. cinerea* known to differ in their aggressiveness on tomato and one strain of *Sclerotinia sclerotiorum*. For this, one-month old plants were subjected to differential fertigation regimes with nitrogen inputs varying from 0.5 to 20 mMol. L⁻¹ NO₃⁻. Plants were grown under those regimes for 3-4 weeks prior to inoculation. Three independent tests were conducted over a three-year period. The fertigation regimes resulted in contrasted development of the lettuce plants and they affected significantly the mineral composition and sugar (fructose, glucose and sucrose) content of the plant tissue, analyzed just before inoculation. Total fresh and dry weights, N, NO₃⁻, P and K contents increased with increasing N concentrations. Leaf area also increased with increasing N concentration up to a threshold level (10 mMol. L⁻¹) above which there was no further increase. In contrast, Ca, Mg and C contents were decreased at higher N levels and the C content did not vary significantly. Contrarily to results obtained in previous studies with tomato, (1) no marked difference in aggressiveness was observed among the six strains, and (2) high N nutrition increased (rather than decreased) the susceptibility of lettuce plants against *B. cinerea*. For all six strains tested, the overall disease severity (assessed by the surface of necrotic lesions), increased with increasing N. The high N fertilization also increased the susceptibility of lettuce to *S. sclerotiorum*. Among all possible factors tested, total N, K and sucrose contents were positively correlated to the surface of the leaf lesions caused by *B. cinerea*, K and sucrose were positively correlated with those caused by *S. sclerotiorum*. Results of this study indicate that increasing NO₃⁻ in the nutrient solution positively affected vegetative growth of lettuce plants but it also increased the susceptibility of plants to *B. cinerea* and *S. sclerotiorum* infection in lettuce.

Keywords: Grey mould, susceptibility, host resistance, N fertilization, *Lactuca sativa*

Journées Jean Chevaugeon 2012



9^{èmes} Rencontres de Phytopathologie-Mycologie de la Société Française de Phytopathologie (SFP)

16 au 20 janvier 2012

Centre Paul Langevin, CAES du CNRS
Aussois (Savoie) – France

<http://colloque.inra.fr//jic2012>



INRA



cirad

LA RECHERCHE AGRONOMIQUE
POUR LE DÉVELOPPEMENT



AMBASSADE DE FRANCE
AU JAPON

SERVICE POUR LA SCIENCE
ET LA TECHNOLOGIE



Science For A Better Life