

Implication of the vacuolar AtPAP26 in the sugars metabolism in Arabidopsis thaliana roots

Antoine Monier, Sarah Monllor, Marie-Hélène Andrieu, Sophie S. Colombie, Frédéric Delmas, Delphine Lapaillerie, Marc Bonneu, Veronique Germain, Martine Dieuaide Noubhani, Bertrand Beauvoit

▶ To cite this version:

Antoine Monier, Sarah Monllor, Marie-Hélène Andrieu, Sophie S. Colombie, Frédéric Delmas, et al.. Implication of the vacuolar AtPAP26 in the sugars metabolism in Arabidopsis thaliana roots. 5. Journées scientifiques du Réseau Français de Métabolomique et Fluxomique, Labo/service de l'auteur, Ville service., May 2011, Paris, France. hal-02749345

HAL Id: hal-02749345 https://hal.inrae.fr/hal-02749345v1

Submitted on 3 Jun 2020 $\,$

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers. L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

F4-P4

Implication of the vacuolar AtPAP26 in the sugars metabolism in Arabidopsis thaliana roots

<u>Antoine Monier¹</u>, Sarah Monlort², Marie-Hélène Andrieu², Sophie Colombier², Frédéric Delmas¹, Delphine Lapaillerie*, Marc Bonneu*, Véronique Germain¹, Martine Dieuaide-Noubhani¹ & Bertrand Beauvoit¹

- 1 Université de Bordeaux, UMR 1332, Biologie du Fruit et Pathologie, 71 av Edouard Bourlaux, 33140 Villenave d'Ornon, France
- 2 INRA UMR 1332, Biologie du Fruit et Pathologie, 71 av Edouard Bourlaux, 33140 Villenave d'Ornon, France
- * Plateforme Protéomique de Génomique Fonctionnelle Bordeaux, 146 rue Léo Saignat, 33000 Bordeaux, France

«Purple Acid Phosphatases» (PAPs) are acid phosphatases found in both the animal and vegetal kingdoms. They are characterized by the presence of seven conserved amino acids involved in a dimetal nuclear center and by their pink/purple color in solution [1]. In plants, some purified PAPs have a phosphatase activity on phosphorylated metabolites (PEP, G6P ...) and peptides (e.g. glycosidases) [2]. Furthermore, a peroxidase activity has demonstrated in vitro [3]. PAPs could be implicated both in the metabolism regulation and in the detoxification of reactive oxygen species [4]. To date, 29 different genes encoding PAPs were identified in the *Arabidopsis thaliana* genome. Some of them are induced by Pi deprivation, oxidative burst or senescence. The gene encoding the protein AtPAP26 (At5g34850) exhibits high sequence similarity with orthologs from other plants [3]. The AtPAP26 protein, showing both a phosphatase activity on PEP and G6P and a peroxidase activity, was localized in the vacuole [5] but also appears to be one of the principal root-secreted acid phosphatases [6].

Our aims are to: 1° / analyse the potential role of AtPAP26 in the sugars metabolism and in their vacuolar storage. Our first hypothesis is that PAP26 could be involved in the glucose accumulation in the vacuole *via* its phosphatase activity [7]. To perform this, we are following the incorporation of U-C¹⁴ labeled hexoses in Arabidopsis roots to measure sugars turnover. 2° / Verify the potential implication of AtPAP26 on the dephosphorylation of target proteins.

Références bibliographiques :

[1] Li et al. Journal of Biological Chemistry. Purple acid phosphatases of Arabidopsis thaliana. Comparative analysis and differential regulation by phosphate deprivation. (2002) 277: 27772-81

[2] Bozzo *et al. European Journal of Biochemistry*. Purification and characterization of two secreted purple acid phosphatase isozymes from phosphate-starved tomato (*Lycopersicon esculentum*) cell cultures. (2002) 269: 6278-86
[3] Veljanovski *et al. Plant Physiology*. Biochemical and molecular characterization of AtPAP26, a vacuolar purple acid phosphatase up-regulated in phosphate-deprived Arabidopsis suspension cells and seedlings. (2006) 142:1282-93
[4] del Pozo *et al. Plant Journal*. A type 5 acid phosphatase gene from Arabidopsis thaliana is induced by phosphate starvation and by some other types of phosphate mobilising/oxidative stress conditions. (1999) 19:579-89

[5] Hurley *et al. Plant Physiology*. The dual-targeted purple acid phosphatase isozyme AtPAP26 is essential for efficient acclimation of Arabidopsis to nutritional phosphate deprivation. (2010) 153:1112-22

[6] Tran *et al. Plant Science*. Feeding hungry plants: The role of purple acid phosphatases in phosphate nutrition. (2010) 179:14-27

[7] Alonso *et al. Plant Physiology*. A new substrate cycle in plants. Evidence for a high glucose-phosphate-to- glucose turnover from in vivo steady-state and pulse-labeling experiments with $[^{13}C]$ glucose and $[^{14}C]$ glucose. (2005) 138:2220-32

Mots-clés : Arabidopsis thaliana, sugar metabolism, roots, vacuole