



HAL
open science

Contribution of microbial core-collections and exploitation of polygenic resistances to the development of Phytophthora control strategies in solanaceous crops

Marie-Line Kuhn, Anne Massire, Melissa Cantet, Rene Damidaux, Elvina Gibowski, Youna Ruellan, Véronique Lefebvre, Franck Panabières

► To cite this version:

Marie-Line Kuhn, Anne Massire, Melissa Cantet, Rene Damidaux, Elvina Gibowski, et al.. Contribution of microbial core-collections and exploitation of polygenic resistances to the development of Phytophthora control strategies in solanaceous crops. Plant Resistance Sustainability 2012 International Conference, Oct 2012, La Colle sur Loup, France. hal-02749550

HAL Id: hal-02749550

<https://hal.inrae.fr/hal-02749550>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

PLANT RESISTANCE SUSTAINABILITY

International Conference 2012



La Colle-Sur-Loup (France)
October 16th-19th, 2012



Scientific Programme and Abstracts

Sessions

Session 1: Impact of plant disease resistance on the structure and evolution of pathogen populations

Session 3: From plant-pathogen molecular interactions to the durability of resistance

Session 2: Sustainable and integrated breeding and deployment of genetic resistance

Session 4: Socio-economic issues related to the use of resistant varieties and their deployment in agro-systems

Invited Speakers

Philippe Baret, Université Catholique de Louvain, Belgium - **James Brown**, John Innes Centre, England - **Marion Desquilbet**, INRA, France - **Sylvain Gandon**, CNRS, France - **Benoit Moury**, INRA, France - **Chris Mundt**, Oregon State University, USA - **Laura Rose**, Heinrich-Heine University, Germany - **Walter Rossing**, Wageningen University, The Netherlands - **Peter Thrall**, CSIRO Plant Industry, Australia

Organised by the Institut National de la Recherche Agronomique (INRA)
Metaprogramme on Sustainable Management of Crop Health (SMaCH)



Contribution of microbial core-collections and exploitation of polygenic resistances to the development of *phytophthora* control strategies in solanaceous crops

Kuhn M-H.^{1*}; Massire A.^{2*}; Cantet M.²; Damidaux R.²; Gibowski E.¹; Ruellan Y.²; Lefebvre V.^{2}; Panabières F.^{1**}**

Marie-Line Kuhn^{1*}, Anne Massire^{2*}, Mélissa Cantet², René Damidaux², Elvina Gibowski¹, Youna Ruellan², Véronique Lefebvre²⁼, Franck Panabières¹⁼

¹UMR INRA/CNRS/UNS 1355 Institut Sophia Agrobiotech, 400 route des Chappes, 06903 Sophia-Antipolis, France;

²INRA, UR 1052 Génétique et Amélioration des Fruits et Légumes, Domaine St Maurice, Allée des Chênes, BP 94, 84143 Montfavet cedex, France.

*contributed equally to this work;

**contributed equally to this work

Abstract

Oomycetes form a diverse group of fungus-like eukaryotic microorganisms that include saprophytes as well as pathogens of animals and some of the most devastating pathogens of dicotyledonous plants. Among them, few species of the genus *Phytophthora*, such as *P. capsici* and *P. infestans* cause enormous economic damage on Solanaceous crops. Major genes have been identified in tomato to confer resistance against *P. infestans*, but were rapidly overcome by virulent populations. Pepper resistance towards *P. capsici* is provided by several quantitative trait loci (QTLs). Efficient exploitation of such polygenic resistance sources in plant breeding programs requires a good evaluation of their spectrum and durability. This requires a precise knowledge of the diversity of pathogen populations (with a focus on virulence, aggressiveness and host range) and some of their adaptive traits, such as changes in virulence or adaptation to prevalent cultivars. Theoretically, the use of *Phytophthora* core-collections reflecting the overall intraspecific diversity would help to assess the *a priori* durability of resistance sources.

We initiated a study to evaluate the potential sustainability of polygenic pepper and tomato resistances to *Phytophthora*. It implied i) to estimate the diversity of *P. infestans* and *P. capsici* collections and their evolution over time; ii) to develop diverse quantitative and qualitative pathogenic assays to identify the various components underlying polygenic resistances. Collections included isolates from various geographic origins. Genetic diversity was investigated using a combination of neutral markers and genes under positive selection. First results on *P. capsici* reveal significant variations in the structure and dynamics of regional populations, and suggest that *P. capsici* core-collections may be of local interest. We also showed that *P. infestans* populations are diverse but structured. Hence, strains isolated on tomato or potato differ at the genotypic and phenotypic levels. The use of *P. infestans* core-collections in breeding programs therefore requires gaining knowledge on the structure and epidemiological characteristics of natural pathogen populations collected on tomato.

Keywords: Resistance, *Phytophthora spp.*, pepper, tomato, core-collection, plant breeding.