



**HAL**  
open science

## Sustainability of three winter cereals based cropping systems (FR)

Clotilde Toque-Rouillon, Jean-François Garnier, Lionel Jouy, Patrick Retaureau, Gabriele Fortino, Valérie Leveau

► **To cite this version:**

Clotilde Toque-Rouillon, Jean-François Garnier, Lionel Jouy, Patrick Retaureau, Gabriele Fortino, et al.. Sustainability of three winter cereals based cropping systems (FR). Future IPM in Europe, Mar 2013, Riva del Garda, Italy. 334 p., 2013, Future IPM in Europe. hal-02749574

**HAL Id: hal-02749574**

**<https://hal.inrae.fr/hal-02749574>**

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.



# SUSTAINABILITY OF THREE WINTER CEREALS BASED CROPPING SYSTEMS (FR)

Toque-Rouillon Clotilde <sup>(1)</sup>, Garnier Jean-François <sup>(1)</sup>, Jouy Lionel <sup>(1)</sup>, Retaureau Patrick <sup>(1)</sup>, Fortino Gabriele <sup>(2)</sup>, Leveau Valérie <sup>(1)</sup>



(1) ARVALIS-Institut du végétal, Département Recherche et Développement - 91720 BOIGNEVILLE (FR)  
 (2) INRA, UAR 1240 Eco-Innov, BP 01 - 78850 THIVERVAL-GRIGNON (FR)

## Description of the 3 Systems :

Description			Crop Rotations	Main IPM Tools
<b>AS</b>	<b>Advanced System</b>	Organic system	AA(x2) – WSW – legumes – S Flax – WFP - WSW	Crop sequence Plough, Stubble cultivation Tolerant cultivars, Alfa-alfa management Mechanical weeding, Delayed sowing dates
<b>IS</b>	<b>Intermediate System</b>	Integrated system	SF – WHW – mustards+legumes – SB- WSW	Crop sequence Stubble cultivation, Plough if necessary Tolerant cultivars, Delayed sowing dates, Low nitrogen levels Mechanical weeding
<b>CS</b>	<b>Current System</b>	Conventional system	WORS – WSW – mustard – SB – WHW	Stubble cultivation, Plough if necessary Productive cultivars septioria-tolerant Fertilisation and crop protection to maximize gross margin


\* WSW = Winter Soft Wheat ; AA = Alfa-Alfa ; S Flax = Spring Flax ; WFP = Winter Faba Pea ; SF = Sun Flower ; WHW = Winter Hard Wheat ; SB = Spring Barley ; WORS = Winter Oil RapeSeed

## The Context :



- **Site :** Boigneville (FR)
- **Soil :** Clayeley-silty soil over chalk
- **Climate :** degraded oceanic
- **Annual rainfall :** 600 mm
- **Field capacity :** 170 mm
- **Economic context :** arable crops systems without livestock
- **Pests :** insects and slugs at a minimal level, disease impact about 1 to 2 tons/year

## Assesment with DEXI-PM Model :



## Example of some Criteria :



Average/hectare for the cropping system	Average 2008 – 2010			Year 2012		
	AS	IS	CS	AS	IS	CS*
TFI – total (ref 5.9)	0	1.6	3.1	0	2.0	4.6
TFI – herbicides (ref 1.7)	0	1	1.7	0	1.4	2.0
Quality of the weed control (/10)	5.5	7.1	8.2	5	7.3	8.5
Mineral N (kg/ha)	0	80	180	0	120	120
WSW yield (t/ha)	3.4	5.9	8.0	3.0	7.3	8.6
Gross margin (€/ha)	1275	767	956	1445	1090	1180
Net margin (€/ha)	536	168	370	610	430	730
Fuel consumption (L/ha)	74	95	87	79	81	81
Tractor time (h/ha)	3.5	4.0	3.7	3.6	3.5	3.4

\* Winter Oil RapeSeed replaced by Spring Peas

The research leading to these results has received funding from the European Union Seventh Framework Programme (FP7/ 2007-2013) under the grant agreement n°265865- PURE

## First Conclusions :

- ✓ **Currently :**
  - Succeeded in implementing the IPM solutions
  - Positive impact on the use of pesticides, but depending on the pressure of the year
  - Close link between pesticides use and productivity
- ✓ **In the future :**
  - Optimising weed control
  - Improve quality of the products
  - Assess the vulnerability of the results to the context
  - Assess the risk of pesticides transfert into groundwater
  - Assess the risk of ecotoxicity

