



Alopecurus myosuroides transcriptome database (ALOMYBase): unravelling the genetic bases of non-target-site-based resistance to herbicides

Jeanne Gardin, Jerome Gouzy, Sebastien Carrere, Roland Beffa, Christophe C. Delye

► To cite this version:

Jeanne Gardin, Jerome Gouzy, Sebastien Carrere, Roland Beffa, Christophe C. Delye. Alopecurus myosuroides transcriptome database (ALOMYBase): unravelling the genetic bases of non-target-site-based resistance to herbicides. 16. European Weed Research Symposium (EWRS), European Weed Research Society (EWRS). GBR., Jun 2013, Samsun, Turkey. pp.259. hal-02749853

HAL Id: hal-02749853

<https://hal.inrae.fr/hal-02749853>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

2013-99

16th SYMPOSIUM

Samsun 2013

24-27 June, 2013



EWRS

*European Weed
Research Society*

PROCEEDINGS

ISBN: 978-90-809789-12

EWRS European Weed Research Society
 Europäische Gesellschaft für Herbolologie
 Société Européenne de Malherbologie

EWRS 16th SYMPOSIUM, Samsun 2013

Copyright The reproduction of these Proceedings is not permitted without the written permission from the European Weed Research Society

Copies of the Proceeding may be obtained from:

EWRS Bookstore
 P.O. Box 28
 NL-6865 ZG Doorwerth
 The Netherlands

Printing: Erol Offset, Samsun

ISBN: 978-90-809789-12

Program Committee

Chairman: Paolo Barberi (President of the EWRS)

Members: Josef Soukup (Scientific Secretary of the EWRS)

László Radics (Vice President of the EWRS)

Session Organizers

H. Kraehmer	Bayer CropScience, AG Industriepark Hoechst Frankfurt am Main Germany
H. Darmency	INRA, UMR1347, Agroécologie, BP86510, 21065 Dijon, France
M. Vurro	Istituto di Scienze delle Produzioni Alimentari CNR, via Amendola, Bari, Italy
I. Bastiaans	Crop and Weed Ecology Group, Dept. of Plant Sciences, Wageningen University, Wageningen, The Netherlands
G. Economou	Faculty of Crop Production Science, Agricultural University of Athens, Iera Odos Str., Athens
E. Pannacci	Department of Agricultural and Environmental Sciences, University of Perugia, Borgo XX Giugno 7 Perugia
B. Melander	Department of Integrated Pest Management, University of Aarhus, Forsøgsvej Slagelse
P. Hatcher	School of Biological Sciences, Plant Science Laboratories, University of Reading, Gb-Reading
P. Kudsk	Department of Integrated Pest Management, University of Aarhus Forsøgsvej 1, Slagelse
N. Uygur	Department of Plant Protection, University of Çukurova, Adana, Turkey
P. Neve	School of Life Sciences, University of Warwick, CV4 7AL, Coventry, United Kingdom
B. Rubin	Faculty of Agricultural, Food and Environmental Sciences, Hebrew University of Jerusalem, RH Smith Inst. Plant Sci. & Genetics, Rehovot
S. Christensen	University of Copenhagen, Faculty of Life Sciences Department of Agriculture and Ecology, Højbakkegård Allé, Taastrup Institute for Land Use, University of Rostock, Satower Str. Rostock

Individual-based simulation as a tool to help understand how ecology, management and genetics affect the evolution of herbicide resistance

M. Renton¹, K. Flower¹, R. Busi¹, D. Thornby², S. Manali¹, G. Gill³, S. Powles¹

¹School of Plant Biology, Australian Herbicide Resistance Initiative and UWA Institute of Agriculture, University of Western Australia, Crawley WA 6009, Australia

²Queensland Department of Agriculture, Fisheries and Forestry, Toowoomba QLD 4350, Australia

³School of Agriculture, Food and Wine, University of Adelaide, North Terrace, Adelaide SA 5005, Australia

michael.renton@uwa.edu.au

Herbicide resistance is a growing problem threatening world food production. The way weeds evolve resistance to herbicides depends on a complex interaction between the underlying genetics, the weed management used by the farmer, and the biology of the weed species. Evaluating different management options for delaying or avoiding resistance evolution is a challenge because it depends on this complex interaction of multiple factors at different scales, and because it is a long-term large-scale process that is difficult to investigate properly with small-scale short-term experiments. Simulation modelling provides a means of integrating the various factors involved in resistance, including physiological mechanisms, molecular and population genetics, ecology, evolutionary dynamics, and practical management. To help better understand, predict and manage evolution of herbicide resistance, we developed a model, known as PERTH (Polygenic Evolution of Resistance To Herbicides), to predict the rates and patterns of resistance evolution in a wide range of different conditions. PERTH is based on an individual-based simulation methodology which allows it to represent a wide range of details of the real biological system. PERTH has been used to help evaluate the efficacy of management options such as herbicide rotation, maintaining robust herbicide rates at high efficacy, avoiding or introducing refuges, and using non-chemical methods of control, and how the efficacy of these options depends on the genetics underlying the resistance. For example, modelling results showed that maintaining robust herbicide rates at high efficacy is an important strategy for avoiding polygenic resistance, but has a negligible effect on the evolution of monogenic resistance. PERTH has also been used to show that the evolution of resistance can affect be affected by ecological functional traits of different weed species, such as seed bank longevity and dormancy, fecundity, seed dispersal, and rates of self-fertilisation versus out-crossing. Recently PERTH has been adapted to account for spatial heterogeneity, which has let us predict that factors such as seed and pollen dispersal can strongly affect the evolution of resistance. We conclude that the strength of simulation models, and of individual-based approaches in particular, is that they provide a means of integrating knowledge from multiple scientific disciplines at a wide range of temporal and spatial scales. Thus they provide a valuable tool to help understand how ecology, management and genetics affect the evolution of herbicide resistance, and ultimately help safeguard the sustainability of world food production.

Alopecurus myosuroides transcriptome database (ALOMYBase): Unravelling the genetic bases of non-target-site-based resistance to herbicides

J. Gardin¹, J. Gouzy², S. Carrere², R. Beffa³, C. Délye¹

¹INRA Dijon, UMR Agroécologie, Dijon, France

²INRA Toulouse, Laboratoire Interactions Plantes Micro-organismes, Castanet Tolosan, France

³Bayer CropScience, Institut für Herbizidforschung, Frankfurt/Main, Germany

jeanne.gardin@dijon.inra.fr

Herbicides are the most effective weed control tools, but their efficacy is jeopardized by the evolution of resistances in weeds. Non-target-site-based resistance (NTSR) is the most challenging aspect of herbicide resistance. NTSR is complex, widespread and unpredictable. It can confer resistance to a range of herbicides with different modes of action. Thus, understanding the genetic bases and the evolution of NTSR is crucial to sustain herbicide efficacy. Therefore, the objective of this study is to identify genes involved in NTSR. NTSR is part of the weed response to the herbicide stress. Accordingly, differences in gene expression between resistant and sensitive plants are considered to be at the root of NTSR. Thus, a transcriptomics-based approach is expected to identify NTSR candidate genes. Such an approach is now feasible with the tremendous progresses achieved by the sequencing technologies (next-generation sequencing). This study focuses on blackgrass (*Alopecurus myosuroides*), a major weed in winter crops in Western Europe that evolved NTSR to the most effective herbicides used to control it (i.e., acetolactate synthase and acetyl-CoAcarboxylase inhibitors).

Using the illumina sequencing technology, an exhaustive transcriptome database was established for *A. myosuroides*. This database was used as a reference to compare the transcriptomes of herbicide-sensitive and herbicide-resistant *A. myosuroides* individuals before and after herbicide treatment along a time-course experiment, using a RNA-Seq approach. Genes showing a constitutive or an herbicide-induced increased expression in resistant plants were sought as candidate NTSR genes. 42 genes fulfilling this criterion were identified. These genes encoded enzymes potentially involved in herbicide degradation or compartmenting, regulator proteins, and putative proteins with unknown functions. The differences in expression observed were confirmed by quantitative real time PCR, yielding a set of candidate NTSR genes of interest. Further investigation on the role of these genes in NTSR must now be conducted.