



Physiological and behavioural adaptation to heat stress in 5-week old chickens.

Christine Leterrier, Yrina Colina, Anne Collin, D. Bastianelli, Paul
Constantin, Vasco de Basilio

► To cite this version:

Christine Leterrier, Yrina Colina, Anne Collin, D. Bastianelli, Paul Constantin, et al.. Physiological and behavioural adaptation to heat stress in 5-week old chickens.. 5. Combined Workshop on Fundamental Physiology and Perinatal Development in Poultry, Aug 2011, Wageningen, Netherlands. hal-02749857

HAL Id: hal-02749857

<https://hal.inrae.fr/hal-02749857>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

5th Combined Workshop

Fundamental Physiology and Perinatal Development in Poultry



August 31 - September 3, 2011

Wageningen, the Netherlands



Physiological or behavioural adaptation to heat stress in 5-week old chickens

Leterrier¹, C., Y. Colina², A. Collin³, D. Bastianelli⁴, P. Constantin¹, and V. De Basilio²

¹INRA, UMR 85 Physiologie de la Reproduction et des Comportements, F- 37380 Nouzilly, France,

²Universidad central del Venezuela, Maracay, Edo-Aragua, Venezuela, ³INRA, UR 83 Recherches Avicoles, F-37380 Nouzilly, France, ⁴CIRAD, Systèmes d'élevage méditerranéens et tropicaux, 34 398 Montpellier cedex 05, France

Adaptation to high temperature increases is improved by thermal stimulations during the first week of age in chickens. The present experiment was designed to look whether an adaptation capacity was still present at higher ages (5 weeks of age). We evaluated adaptation by measuring panting behaviour and body temperature during repeated rises in room temperature.

When the chickens were 35 days old, the room temperature (22°C) was increased from 22°C up to 27°C for 2H30, then up to 32°C for 2H30 more, and then switched back to 22°C. On Day 36 and 37, room temperature was increased to 27°C for 6H per day to maintain the thermic stimulation. At Day 38, the same increase in ambient temperature as at Day 35 was repeated. Panting behaviour and body temperature were measured on Day 35 and 38. Body temperature was measured manually and with intra-abdominal telemetric transmitters.

Body temperature increased when the room temperature was set to 27°C and then remained stable during the 2H30. At a room temperature of 32°C, it kept increasing and reached 43.0°C. An adaptation of the thermoregulatory process was found since body temperature at 27°C and 32°C was lower during the second heat challenge (Day 38) compared to the first one (Day 35). Panting frequency was highest at 32°C, but remained unable to maintain body temperature. No difference was recorded in panting behaviour between Day 38 and 35.

Our results suggest that the lower body temperature increase in Day 38 compared to Day 35 would be related to physiological changes rather than changes in animal behaviour.

Keywords: heat stress, adaptation, behaviour, chicken, telemetry

Corresponding author: Dr. Christine Leterrier, INRA, UMR 85 Physiologie de la Reproduction et des Comportements, F- 37380 Nouzilly, France

Email: Christine.Leterrier@tours.inra.fr