



A combination of SNPS is associated with the postprandial chylomicron triacylglycerol response in healthy male adults

Charles Desmarchelier, Jean-Charles Martin, Richard Planells, Marguerite Gastaldi, Marion Nowicki, Aurélie Goncalves, René Valéro, Denis Lairon, Patrick Borel

► To cite this version:

Charles Desmarchelier, Jean-Charles Martin, Richard Planells, Marguerite Gastaldi, Marion Nowicki, et al.. A combination of SNPS is associated with the postprandial chylomicron triacylglycerol response in healthy male adults. 10. International Congress on Coronary Artery Disease, Oct 2013, Florence, Italy. hal-02749985

HAL Id: hal-02749985

<https://hal.inrae.fr/hal-02749985>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

LIPIDS, LIPOPROTEINS AND ATHEROSCLEROSIS

A COMBINATION OF SNPS IS ASSOCIATED WITH THE POSTPRANDIAL CHYLOMICRON TRIACYLGLYCEROL RESPONSE IN HEALTHY MALE ADULTS

C. Desmarchelier¹, J.C. Martin¹, R. Planells¹, M. Gastaldi¹, M. Nowicki¹, A. Goncalves¹, R. Valéro¹, D. Lairon¹, P. Borel¹

¹"Nutrition Obésité et Risque Thrombotique", UMR 1062 INSERM/1260 INRA/ Université d'Aix-Marseille, Marseille cedex 05, France

The postprandial chylomicron (CM) triacylglycerol (TG) response to dietary fat, which is positively associated with atherosclerosis and cardiovascular disease risk, displays a high inter-individual variability. This is assumed to be due, at least partly, to polymorphisms in genes involved in lipid metabolism. Existing studies have so far focused on single SNPs, resulting in a low explained variability. This study aimed to identify a combination of SNPs that could predict an elevated postprandial CM TG response.

Thirty-four healthy male volunteers were subjected to four standardized fat tolerance test meals and genotyped using whole-genome microarrays. Plasma CM TG were measured at regular interval times after each meal. The association of SNPs, in or near candidate genes (126 genes representing 6099 SNPs), with the high (> mean response + 1 SD) vs normal postprandial CM TG response (0 to 8 h postprandial AUC averaged for the four test meals) was assessed by partial least squares discriminant analysis, a multivariate statistical approach.

Data obtained allowed us to generate a validated significant model ($P=2.6.10^{-7}$) that included 28 SNPs in 14 genes (*ABCA1*, *APOB*, *BET1*, *CILP2*, *COBLL1*, *GALNT2*, *IRS1*, *LDLR*, *LIPC*, *MAP3K1*, *MC4R*, *PARK2*, *SLC27A6*, *ZNF664*) and explained 81% of the variance. Univariate analysis revealed several significant ($P<0.05$) differences in CM TG responses between subjects who bore different genotypes in the abovementioned SNPs.

Our results enable us to propose a genetic score that could suggest whether a subject belongs to the high responder phenotype and hence might be at increased risk of atherosclerosis and cardiovascular diseases.

No conflict of interest