



HAL
open science

Glucagon-like peptide 2 (GLP-2) stimulates intestinal function during weaning

T. Thymann, Isabelle Luron Le Huërou-Luron, Y.M. Petersen, M.S. Hedemann, J. Elnif, B.B. Jensen, B. Hartmann, J.J. Holst, P.T. Sangild

► **To cite this version:**

T. Thymann, Isabelle Luron Le Huërou-Luron, Y.M. Petersen, M.S. Hedemann, J. Elnif, et al.. Glucagon-like peptide 2 (GLP-2) stimulates intestinal function during weaning. 44. Annual Meeting of The European Society of Paediatric Gastroenterology, Hepatology and Nutrition (ESPGHAN), May 2011, Sorrento, Italy. hal-02750048

HAL Id: hal-02750048

<https://hal.inrae.fr/hal-02750048v1>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**The 44th Annual Meeting of The European
Society of Paediatric Gastroenterology,
Hepatology and Nutrition**

**25 – 28 May 2011
Sorrento, Italy**

PO-N-0228/PD-N-0126

Pediatric Nutrition

Glucagon-Like Peptide 2 (GLP-2) Stimulates Intestinal Function During Weaning

T. Thymann^{1,*}, I. Le Huërou-Luron², Y. M. Petersen¹, M. S. Hedemann³, J. Elnif⁴, B. B. Jensen⁵, B. Hartmann⁶, J. J. Holst⁶, P. T. Sangild¹

¹Human Nutrition, University of Copenhagen, Frederiksberg, Denmark, ²Unité Mixte de Recherche INRA-Agrocampus, Animal and Human Nutrition, St-Gilles, France, ³Dept. of Animal Nutrition and Physiology, Faculty of Agricultural Sciences, Foulum, ⁴Animal Nutrition, University of Copenhagen, Frederiksberg, ⁵Department of Animal Health and Bioscience, Faculty of Agricultural Sciences, Foulum, ⁶Dept. of Medical Physiology, University of Copenhagen, Copenhagen, Denmark

Objectives and Study: Transition from milk to solid food can be associated with intestinal atrophy and malfunction. The gut hormone, glucagon-like peptide 2 (GLP-2) stimulates gut adaptation. We hypothesized that post-weaning gut adaptation is improved by GLP-2 administration, and that GLP-2 effects vary according to differences in disease sensitivities (e.g. sanitary environment). We used piglets as models for infants.

Methods: In Exp. 1, three week old pigs were weaned in a high-sanitary environment and injected with native GLP-2 (80 µg/kg/12h, n=8) or saline (control, n=8) and compared with preweaning pigs (n=6). In Exp. 2, pigs were weaned in a low-sanitary environment and injected with native GLP-2 (150 µg/kg/12h, n=11) or saline (control, n=11) and compared with preweaning pigs (n=8). In Exp. 3, pigs were weaned in a low sanitary environment and injected with a stabilized acylated GLP-2 analogue (25 µg/kg/12h, n=8) or saline (control, n=8).

Results: Pigs injected with acylated GLP-2 (Exp. 3) showed a lower diarrhea score (2.3 vs. 3.5, $P<0.01$), higher intestinal weight, villi and crypts in the proximal intestine (+20-25%, $P<0.01$) and marked increases in 6 digestive enzymes (+50-100%, all $P<0.05$). Native GLP-2 used in low-sanitary conditions (Exp. 2) increased the density of goblet cells ($P<0.05$), reduced colonic short chain fatty acid levels ($P<0.01$) but did not prevent weaning diarrhea. In the absence of diarrhea in high sanitary conditions (Exp. 1), native GLP-2 injections did not improve gut function. Relative to suckling pigs, intestinal atrophy was similar between controls and GLP-2 treated pigs in both Exp. 1 and 2. In the high-sanitary environment (Exp. 1), the activities of 5 brush border enzymes were markedly increased in both GLP-2 and controls compared with suckling pigs (>300 %, $P<0.05$). In contrast, activities of 3 peptidases were markedly reduced in both GLP-2 and controls in the low-sanitary environment (Exp. 2, -50%, $P<0.001$) compared with suckling pigs. Plasma GLP-2 levels were highest in Exp. 3 (constant levels of ~20 nM), lower in Exp. 2 (daily cycles of 5-30 nM) and lowest in Exp. 1 (daily cycles <1.0 nM).

Conclusion: Exogenous GLP-2 treatment improves gut digestive function during the dietary transition from milk to solid food, but the effects are most pronounced with high-dose GLP-2 and high disease sensitivity. Improved digestive function toward adult type diets may play an important role in the prevention of diarrhea, which is a common clinical response to the weaning transition in both infants and animals.

Disclosure of Interest: None Declared