



**HAL**  
open science

## **Sow influence on neonatal survival: a special focus on colostrum**

Hélène Quesnel, Florence Gondret, Elodie Merlot, Florence Loisel, Chantal Farmer

► **To cite this version:**

Hélène Quesnel, Florence Gondret, Elodie Merlot, Florence Loisel, Chantal Farmer. Sow influence on neonatal survival: a special focus on colostrum. 9. International Conference on Pig Reproduction (ICPR), Jun 2013, Olsztyn, Poland. hal-02750121

**HAL Id: hal-02750121**

**<https://hal.inrae.fr/hal-02750121>**

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## 2. PREGNANCY, PARTURITION AND THE NEONATE

### 2.1. Plenary Session

PP5

## Sow influence on neonatal survival: a special focus on colostrum

Hélène Quesnel<sup>1,2</sup>, Florence Gondret<sup>1,2</sup>, Elodie Merlot<sup>1,2</sup>, Florence Loisel<sup>1,2</sup> and Chantal Farmer<sup>3</sup>

<sup>1</sup>INRA, UMR 1348 PEGASE, F-35590 Saint-Gilles, France; <sup>2</sup>Agrocampus Ouest, UMR 1348 PEGASE, F-35000 Rennes, France; <sup>3</sup>Agriculture and Agri-Food Canada, Dairy and Swine R & D Centre, Sherbrooke, Canada

The main cause of early postnatal deaths in piglets is hypothermia due to an inadequate intake of colostrum. Colostrum consumption is the outcome of complex interactions between the sow, the piglet, the litter and the environment. The sow may have an impact on many factors that are determinant for colostrum intake and chances of survival, such as piglet weight, maturity and vitality at birth, or within-litter variation in birth weight. Colostrum intake also depends on the ability of the sow to produce colostrum in sufficient quantity to fulfill the needs of the whole litter. Maternal stress during gestation may increase piglet morbidity and mortality up to weaning, presumably by affecting the ontogeny of the fetal immune system, but also IgG contents in colostrum and IgG transfer to newborn piglets. Ways to reduce neonatal mortality through maternal feeding during gestation are largely investigated. Feeding strategies generally failed to increase piglet birth weight but led to more promising results on piglet maturity and vitality at birth and on the acquisition of passive immunity. There is some evidence that maternal feeding during the periparturient period may influence both the quantity and the quality of colostrum; this needs however to receive further attention.