

Genes flow in populations of *Bremia lactucae*, the causal agent of downy mildew of lettuce, in France

Romain Valade, Brigitte Maisonneuve, Claire Neema

► **To cite this version:**

Romain Valade, Brigitte Maisonneuve, Claire Neema. Genes flow in populations of *Bremia lactucae*, the causal agent of downy mildew of lettuce, in France. Congrès du réseau de Génétique Moléculaire des Oomycètes, Jun 2010, Toulouse, France. hal-02750444

HAL Id: hal-02750444

<https://hal.inrae.fr/hal-02750444>


Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Congrès du Réseau de Génétique
Moléculaire des Oomycètes
6-8/6/2010 - TOULOUSE**

**OOMYCETE MOLECULAR GENETICS
NETWORK CONGRESS**



Congress
of the Oomycete Molecular Genetics
Network

Congrès
du Réseau de Génétique Moléculaire
des Oomycètes

6-8/6/2010
Salons Marengo
Médiathèque José Cabanis,
Toulouse, France

Genes flow in populations of *Bremia lactucae*, the causal agent of downy mildew of lettuce, in France

VALADE R.¹, MAISONNEUVE C.², AND C. NEEMA¹

¹*AgroParisTech, UMR BIOGER-CPP, Thiverval Grignon, 78850, France*

²*INRA, GAFL Domaine St Maurice, 84143 Montfavet cedex, France*

Bremia lactucae, the causal agent of downy mildew on *Lactuca sativa*, is an important economic problem for lettuce crops. Breeders used genetic control to limit damages of this pathogen and more particularly specific resistances. However, under selection pressure, *B. lactucae* populations showed a rapid adaption to host resistances. A better knowledge is necessary to understand this rapid virulence evolution. A study was initiated to evaluate genetic diversity of this biotrophic, heterothallic and diploid pathogen, and to determine the impact of recombination and migration on the structure of pathogen populations. More than four hundred isolates were recovered in France from different regions and different varieties (carrying different resistance genes) from *L. sativa* and *L. serriola* which is a wild host species. More than one hundred isolates were characterized for their virulence. Neutral markers, microsatellites, were developed to assess population structure and gene flow in french populations of *Bremia*. Moreover, polymorphism in RxLR candidate genes was evaluated in relation with genotypic and phenotypic diversity. Results showed an important polymorphism for virulence in *B. lactucae* french populations and the occurrence of gene flow between populations and between the wild and crop pathosystems. These results will be discussed regarding durable resistance management.