



Demonstrating the functional role of traits for plant performance: a mechanistic modeling approach.

Nicolas Gross, Vincent V. Maire, Raphaël Martin, Jean-François J.-F. Soussana

► To cite this version:

Nicolas Gross, Vincent V. Maire, Raphaël Martin, Jean-François J.-F. Soussana. Demonstrating the functional role of traits for plant performance: a mechanistic modeling approach.. British Ecological Society, annual Meeting, British Ecological Society (BES). GBR., Sep 2008, Londres, United Kingdom. 20 p. hal-02750769

HAL Id: hal-02750769

<https://hal.inrae.fr/hal-02750769>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

*Demonstrating the functional role of
traits for plant performance:
a mechanistic modeling approach*

Gross N, Maire V, Hill D, Martin R, Soussana JF



What are plant traits?



A simple definition:

Morphological or physiological characteristics with a function for plant to subsist in its local environment


→ Response traits (Lavorel et al. 2007 for review)

Traits co-vary along axis of specialization:

Notion of trade-off and trait syndrome

Conservative syndrome


Exploitative syndrome



Diaz et al. 2004

Traits co-vary along axis of specialization:

Notion of trade-off and trait syndrome



Difficulty to test effect of particular trait
on plant performance due to trade-off among traits

A modeling approach to isolate trait effect on plant performance

GEMINI

(Grassland Ecosystem Model with INdividual centered Interactions)

Environmental variables

Demographic
models

Ecophysiologic
biochemistry
models

Demography
&
Morphology


*Population Dynamics
Level of biodiversity*

*Individual centred
Model*

Flux Models
C, N, H₂O

*Acquisition
and
Utilisation*





GEMINI is composed by different submodels

Overall 74 physiological and morphological parameters are needed for plants

Parameterized and evaluated on 13 grasses species (monoculture) grown in 2003-2004
Maire et al. in prep

More informations see: www2.clermont.inra.fr/discover/gemini/model.htm
 & www.the-jena-experiment.de

Testing traits effect on plant performance using a sensitivity analysis

4 traits were selected

two leaf traits (LDMC, Ph0) and two stature traits (D0, L0)

Hypothesis

Strong linkages within leaf traits (D0 and L0) and within stature traits (LDMC, Ph0) (Wright et al. 2004)

Independence between Leaf traits and stature traits (Ackerly 2004, Gross et al. 2007)

Testing traits effect on plant performance using a sensitivity analysis

12 grasses species were used


Tested only effect of morphological traits
 Monoculture at equilibrium
 No Reproductive stage
 No water limitation
 Environnement 2003-2004

Traits space exploration

10 value of traits per species
 in a full factorials design

In total


$12 \text{ species} \times 10 \text{ value}^4 \text{ traits} \times 2 \text{ N conditions} = 240\,000$
 simulations ran with monoculture growth conditions 2003/2004





Trade-off at inter-specific level



Plant biomass in function of traits value (Plant Height)
all other parameter are fixed



One trait value maximize plant performance

In the 4D space traits:

traits combinations which maximize plant performance



(variance explained >80 %)

Traits combinations which maximize plant performance are in accordance
Trade-off among traits and among species is well predicted by the model
which current plant schema (Grime 1977, Westoby 1998)

Some insight coming from the modeling approach

→ Plant traits observed in the field maximize resource utilization and plant performance

Best light interception, N absorption, N leaf concentration predicted in the

model

→ Each species can be view has an island, a particular traits combination which locally maximize plant fitness in the traits space

→ Our study did not show why species differed in traits syndrome (Selection along ecological gradients...), but can explain why trade-offs among traits exist (all combinations are not equal)






Trade-off at intra-specific level

and interaction among traits

Interaction between tiller density and plant height



Ridge which maximize plant performance

Consistently with our hypothesis there is a negative trade-off between tiller density and plant height

Constraint between traits change depending on plant syndrome

Conclusion

- By using a modeling approach we demonstrated the functional role of traits for plant performance
- Species are organized along trade-off of traits where species maximize their performance
- Trade-off exist both between and within species but are not necessarily the same

Perspective


Studying the impact of plant plasticity in response to change in abiotic conditions

Linkage between species interactions and plant traits: Plasticity, convergence, divergence...



Thank you!!!

Interaction between tiller density and LDMC




Within species there is a strong linkage between LDMC and tiller density

Not necessarily observed at inter-specific level

Constraint between traits change depending on plant syndrome

Some combinations in the traits space are equivalent



→ Not same level of plasticity in function of traits syndromes







Interactions between two traits: Four possibilities


Biomass



Biomass



In the 4D space traits: traits combinations which maximize plant performance



This ACP is concordant with the Grime Model GSR (1977).
 Traits maximized by the model are strongly related with observed traits measured in the field