

# Genetic and molecular diversity of fruit traits in melon

Catherine Dogimont

► **To cite this version:**

Catherine Dogimont. Genetic and molecular diversity of fruit traits in melon. Indo French Symposium, Jan 2010, Lucknow, India. hal-02750884

**HAL Id: hal-02750884**

**<https://hal.inrae.fr/hal-02750884>**

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

# INDO FRENCH SYMPOSIUM

## GENOMICS AND BIOTECHNOLOGY OF FRUIT QUALITY: RECENT ADVANCES

January 18-20, 2010, Lucknow, India



 Indo French Council for Promotion of Advance Research, India



National Botanical Research Institute, Lucknow, India



Ecole Nationale Supérieure Agronomique Toulouse, Tolosan, France

## **GENETIC AND MOLECULAR DIVERSITY OF FRUIT TRAITS IN MELON**

**Catherine DOGIMONT**

INRA Avignon, UR1052, Unité de Génétique et Amélioration des Fruits et Légumes,  
Domaine St Maurice BP 94- 84143 Montfavet cedex France

Melon is an important horticultural crop across wide areas of the world. Different uses and diverse consumption habits lead to the wide diversity of fruit types - sweet and non-sweet melon types- and to a large morphological variation in fruit characteristics. Thus, interestingly, both climacteric and non-climacteric fruit ripening exist in melon. We investigated the genetic variation of the melon species using a large collection of accessions maintained at INRA and studied the genetic control of fruit traits. In order to shed light on the molecular processes underlying fruit ripening in melon, several genes involved in the ripening process were identified by sequence homology to tomato or Arabidopsis and were mapped on the melon genome. A reverse genetic approach was used to assign a functional role to these genes in melon. EMS mutants obtained from a climacteric melon of the Charentais type were selected by TILLING. Phenotyping of the mutants is ongoing. Using an EcoTILLING approach, we are investigating the allelic diversity of the genes, available within a melon collection.