



Monitoring the rumen microbial populations changes during feed-induced lactic, butyric or propionic acidosis in sheep

Abderzak Lettat, Pierre Noziere, Mathieu Silberberg, Diego Morgavi, Cécile Martin

► To cite this version:

Abderzak Lettat, Pierre Noziere, Mathieu Silberberg, Diego Morgavi, Cécile Martin. Monitoring the rumen microbial populations changes during feed-induced lactic, butyric or propionic acidosis in sheep. 7. Joint Symposium of Rowett-INRA 2010, Jun 2010, Aberdeen, United Kingdom. hal-02751115

HAL Id: hal-02751115

<https://hal.inrae.fr/hal-02751115>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.




**Rowett Institute
of Nutrition and Health**
University of Aberdeen



Rowett-INRA 2010

23-25 June 2010

Aberdeen Exhibition and Conference Centre



Gut Microbiology:

new insights into
gut microbial ecosystems

7th Joint Symposium organised by the **Rowett Institute of Nutrition and Health**, University of Aberdeen, Scotland (UK) & the **Institut National de la Recherche Agronomique**, Clermont-Ferrand-Theix (France)

Effect of selected antibiotics on growth and metabolism of *Megasphaera elsdenii* NCIMB 41125, a lactate utilising probiotic for ruminants.

Pieter Henning¹, Anu Kettunen², Juha Apajalahti², Frans Hagg¹, Charles Horn¹

¹MS Biotech, Centurion, South Africa, ²Alimetrix Ltd, Helsinki, Finland

Ruminants fed high-concentrate diets run the risk of rumen acidosis, a disorder characterised by lactic acid accumulation and lowered rumen pH. Accumulation of lactic acid can be inhibited by increasing the number and activity of lactic acid utilising bacteria, as demonstrated in previous studies with *Megasphaera elsdenii* NCIMB 41125 (MeCH4). Use of this organism as a probiotic requires certainty that it is not inhibited by prophylactic antibiotics still included in ruminant feeds in many countries outside the EU. The objective of this study was to look at the interaction between various economic important in-feed antibiotics and MeCH4.

An in vitro batch fermentation model using 1 gram of substrate DM (50% corn silage, 25% barley meal, 25% soy meal) in 40 ml buffered rumen digesta was used.

Anaerobic incubation lasted 9 hours at 37°C. Each antimicrobial compound tested was incubated with or without addition of MeCH4, and was added at two levels, selected to give 15 and 35% inhibition of total gas production, respectively. Zinc Bacitracin (0.1 and 0.5mg), Flavomycin (2 and 5mg), Virginiamycin (0.005 and 0.015mg) and Tylosin (0.2 and 1.0 mg) were tested (amount per 40 ml incubation in parentheses). Gas production kinetics, pH, short chain fatty acids, total bacteria and *M. elsdenii* were measured.

MeCH4 greatly reduced ($P < 0.001$ in all cases) the residual level of lactic acid both in the presence and absence of antibiotics. All antibiotics reduced overall bacterial activity, however, addition of MeCH4 significantly ($P < 0.05$) increased activity, as measured by gas production, in each case.

Monitoring the rumen microbial populations changes during feed-induced lactic, butyric or propionic acidosis in sheep

Abderzak Lettat^{1,2}, Pierre Nozière¹, Mathieu Silberberg¹, Diégo P. Morgavi¹, Cécile Martin¹

¹INRA, UR1213 Herbivores, Clermont Ferrand - Theix, France, ²Danisco SAS, Dangé St Romain, France

Ruminal acidosis is a recurrent problem in ruminant production systems. The prevailing fermentation acid is difficult to predict in practice and it is thought to be the reason for the variable effectiveness of some rumen modifiers. We developed a sheep model of experimentally-induced lactic, butyric or propionic acidosis to describe the associated changes in the rumen microbiota. Twelve rumen-cannulated sheep separated into 3 groups of 4 animals were fed a hay:concentrate diet (80:20) during 3 weeks followed by three consecutive days of acidosis challenge by direct intraruminal supply (1.2% BW) of ground wheat, maize or beet pulp. Rumen contents were analysed for fermentation parameters and polysaccharidase activities. Populations of *Fibrobacter succinogenes*, *Streptococcus bovis*, *Lactobacilli* and *Prevotella* were quantified by qPCR.

At the end of the challenge, rumen pH was lower for wheat (4.87) than for maize (5.17) and beet pulp (5.55). Lactate was predominant for wheat (33.8 mM vs. ~3 mM for other challenges), whereas beet pulp and maize favoured a high molar percentage of propionate (22.5%) and butyrate (16.9%), respectively. Fibrolytic activity and *F. succinogenes* proportion were similar among treatments. Amylase activity was highest for wheat, intermediate for maize and lowest for beet pulp. *Lactobacilli* and *S. bovis* proportions tended to be highest for wheat (1.74 vs. ~0.12%; $P = 0.15$) and maize (0.49 vs. ~0.05%; $P = 0.17$) respectively, whereas *Prevotella* was highest for beet pulp (49 vs. ~29%). This study demonstrated that, feed-induced lactic, butyric or propionic acidosis were associated with a specific change in the bacterial population.