



**HAL**  
open science

## Effect of photoperiod on aerial morphogenesis of eight Medicago truncatula lines

Bernadette Julier, Jean-Baptiste Pierre, Philippe P. Barre, Christian C.  
Huyghe

► **To cite this version:**

Bernadette Julier, Jean-Baptiste Pierre, Philippe P. Barre, Christian C. Huyghe. Effect of photoperiod on aerial morphogenesis of eight *Medicago truncatula* lines. 41. North American Alfalfa Improvement Conference & 20. Trifolium Conference, National Alfalfa & Forage Alliance (NAFA). St. Paul, USA., Jun 2008, Dallas, United States. hal-02751436

**HAL Id: hal-02751436**

**<https://hal.inrae.fr/hal-02751436>**

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.


L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

## Effect of photoperiod on aerial morphogenesis of eight *Medicago truncatula* lines

**Bernadette Julier, Jean-Baptiste Pierre, Philippe Barre and Christian Huyghe**  
INRA, UR4, Unité de Recherche Pluridisciplinaire Prairies et Plantes  
Fourragères, 86600 Lusignan, France. Email: [bjulier@lusignan.inra.fr](mailto:bjulier@lusignan.inra.fr)

The influence of environmental factors on aerial morphogenesis is only partly described on the model legume *Medicago truncatula*. Such factors can affect phenotypic evaluation of genetic material such as recombinant inbred lines (RILs), mutants or natural accessions. The objective of this study was to describe the effects of photoperiod, genotype and the genotype x photoperiod interaction on morphogenesis. In growth chambers, six parental lines of RILs populations and two Israeli accessions were grown under two photoperiods, 12 and 18h applied without significant change in total energy supply. Primary branch elongation was measured twice a week, flowering time was scored, and when stem elongation had ceased, the number of primary branches was counted and the length of the main stem was measured.

For all traits, photoperiod and genotype effects were highly significant in analysis of variance but the effect of photoperiod was the highest. On average, plants flowered earlier and had longer main stem, shorter branch length and higher branch elongation rate at a long than at a short photoperiod. Duration of branch elongation and number of branches were lower at a long than at a short photoperiod. The interaction between genotype and photoperiod was always significant. Flowering of Jemalong6 and F83005.5 was more hastened by long photoperiod than that of the other genotypes. Branch length was shorter at a long than at a short photoperiod for five genotypes, longer for one genotype and little affected for two genotypes.



These results show that plant morphology is greatly affected by environmental conditions. A large genetic variation is available. Moreover, the interaction between conditions and genotypes is large, modifying the ranking of the genotypes. It indicates that the protocols for evaluation of RILs, mutants or natural accessions must be carefully chosen.

### **Acknowledgements**

We thank Dr A. Samach from University of Jerusalem (Israel), Dr T. Huguet from INRA-CNRS at Toulouse (France) and J.M. Prospero from INRA at Montpellier (France) for providing the seeds.