



HAL
open science

Linkage disequilibrium mapping for tomato fruit quality traits

Mathilde M. Causse, Nicolas N. Ranc, Stephane Munos, Sophie Rolland, Marie-Christine Le Paslier, Sylvain S. Santoni, Jean-Paul Bouchet, Dominique D. Brunel

► To cite this version:

Mathilde M. Causse, Nicolas N. Ranc, Stephane Munos, Sophie Rolland, Marie-Christine Le Paslier, et al.. Linkage disequilibrium mapping for tomato fruit quality traits. 5. Solanaceae Genome Workshop, Oct 2008, Cologne, Germany. hal-02751766

HAL Id: hal-02751766

<https://hal.inrae.fr/hal-02751766>

Submitted on 3 Jun 2020

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

#158: Linkage Disequilibrium mapping for tomato fruit quality traits

M. Causse¹, N. Ranc², S. Munos², S. Rolland², M. Le Paslier³, S. Santoni⁴, J. Bouchet¹,
D. Brunel⁵

¹INRA - GAFL, Montfavet, France

²INRA, Montfavet, France

³INRA UR EPGV, Evry, France

⁴INRA UMR DIAPC, Montpellier, France

⁵INRA EPGV, Evry, France

Linkage Disequilibrium (LD) mapping links the natural variability present in germplasm collections to molecular polymorphism and allows the identification of candidate genes. We assessed the possibilities of such approach for the study of candidate genes for tomato fruit quality. Knowledge of the genetic diversity and structure of the germplasm is necessary before implementing LD mapping. We genotyped 360 tomato accessions with 20 SSR markers for highlighting the genetic structure of the sample. A genetic continuum between cultivated tomato and wild ancestor *S. pimpinellifolium* was established by a part of *S. lycopersicum* var. *cerasiforme* accessions. The other part of cherry tomato accessions was closer to cultivated accessions. Nested core collections from 8 to 64 accessions of *S. lycopersicum* var. *cerasiforme* were built using molecular and phenotypic data. Cherry tomato accessions were tested for association genetics because (i) they exhibit a large range of variation for fruit quality traits, (ii) they show valuable amount of genetic diversity and (iii) their admixed position between wild and cultivated forms is supposed to increase mapping resolution. Fifty gene fragments from chromosome 2 were sequenced on the largest core collection and SNPs were recorded. LD decreases slowly with genetic distance but the *S. lycopersicum* var. *cerasiforme* admixture group showed reliable results for association genetics even when a limited number of accessions is used. Association could thus be used either to find or to validate candidate genes. SNPs were identified by sequencing 27 fragments of candidate genes for quality traits on the smallest core collection incremented with accessions from *S. lycopersicum* and *S. pimpinellifolium*. SNPs were much more frequent in *S. lycopersicum* var. *cerasiforme* (1 SNPs every 220 bp) than *S. lycopersicum* accessions (1 SNPs every 1000 bp). These SNPs will be used in future studies to genotype the entire collection.