



**HAL**  
open science

## Re-evaluation of the *Globodera pallida* resistance QTLs originating from the wild potato species *Solanum spgazzinii*

Bernard Caromel, Marie-Claire Kerlan, Sarah Danan, Didier Mugniery,  
Véronique Lefebvre

### ► To cite this version:

Bernard Caromel, Marie-Claire Kerlan, Sarah Danan, Didier Mugniery, Véronique Lefebvre. Re-evaluation of the *Globodera pallida* resistance QTLs originating from the wild potato species *Solanum spgazzinii*. 3. Annual Meeting, COST 872 Workshop & MC, May 2009, Toledo, Spain. hal-02751896

**HAL Id: hal-02751896**

**<https://hal.inrae.fr/hal-02751896v1>**

Submitted on 3 Jun 2020

**HAL** is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

**Re-evaluation of the *Globodera pallida* resistance QTLs originating from the wild potato species *Solanum spgazzinii*.**

B. Caromel<sup>1</sup>, M.C. Kerlan<sup>2</sup>, S. Danan<sup>1</sup>, D. Mugniéry<sup>3</sup> & V. Lefebvre<sup>1</sup>

1. INRA-UR1052, Génétique et Amélioration des Fruits et Légumes, BP 94, 84143 Montfavet, FRANCE.
2. INRA - Agrocampus Ouest - Univ Rennes 1, UMR118, Amélioration des Plantes et Biotechnologies Végétales, 29260 Ploudaniel, FRANCE.
3. INRA, Agrocampus Rennes, Univ Rennes 1, UMR1099 BiO3P (Biologie des Organismes et des Populations appliquée à la Protection des Plantes), 35653 Le Rheu, FRANCE

Most of the resistances to the cyst nematode *Globodera pallida*, uncovered in the potato gene pool, originate from wild species related to *Solanum tuberosum*. Using the Interval Mapping (IM) method, we mapped, in a previous study, three QTLs (Quantitative Trait Loci) affecting the number of *G. pallida* newly formed cysts, on the *S. spgazzinii* genome (Caromel *et al.* 2003). Using the Composite Interval Mapping (CIM) method and taking into account the root system size, we re-analysed our data and found that i) one of the previously detected minor-effect QTLs was in fact due to its effect on the variation of root size, ii) it exists minor-effect QTLs originating from the cultivated susceptible parent, which had remained undetected with the IM method, and iii) the individual effects of the previously detected QTLs were overestimated. Evaluation of *G. pallida* development in roots of plants carrying different allelic combinations at resistance QTLs allowed us to confirm or infirm the effect of these QTLs on resistance to *G. pallida*. We conclude that a high level of resistance to *G. pallida* could be obtained by introgressing the *GpaV<sub>spg</sub>* major-effect QTL (formerly called *GpaMI*) and the *GpaXII<sub>spg</sub>* minor-effect QTL (formerly *GpaM3*), originating from *S. spgazzinii*, in a *S. tuberosum* background carrying the *GpaVIII<sub>tr</sub>* QTL.



# **NEMAGENICS**

Exploiting genomics to understand  
plant-nematode interactions

**Proceedings of the Third Annual Meeting**

**Toledo, Spain  
25-28 May 2009**

Organised by



**CSIC**