



HAL
open science

Prediction of forage rumen protein degradability from an in situ degradability database

Marie-Odile M.-O. Nozieres, J Pierre Dulphy, Jean-Louis J.-L. Peyraud, René
Baumont

► To cite this version:

Marie-Odile M.-O. Nozieres, J Pierre Dulphy, Jean-Louis J.-L. Peyraud, René Baumont. Prediction of forage rumen protein degradability from an in situ degradability database. 21. General Meeting of the European Grassland Federation (EGF), Apr 2006, Badajoz, Spain. Coja de badajoz, Grassland Science in Europe, 11, 2006, Sustainable Grassland Productivity. Proceedings of the 21st General Meeting of the European Grassland Federation. hal-02752379

HAL Id: hal-02752379

<https://hal.inrae.fr/hal-02752379>

Submitted on 25 Jan 2023

HAL is a multi-disciplinary open access archive for the deposit and dissemination of scientific research documents, whether they are published or not. The documents may come from teaching and research institutions in France or abroad, or from public or private research centers.

L'archive ouverte pluridisciplinaire **HAL**, est destinée au dépôt et à la diffusion de documents scientifiques de niveau recherche, publiés ou non, émanant des établissements d'enseignement et de recherche français ou étrangers, des laboratoires publics ou privés.

Prediction of forage rumen protein degradability from an *in situ* degradability database



Marie-Odile Nozières ¹, Jean-Pierre Dulphy ¹, Jean-Louis Peyraud ², René Baumont ¹

¹ INRA - Centre de Clermont-Ferrand / Theix, 63122 St-Genès-Champanelle, France

² INRA – Centre de Rennes, 35590 St-Gilles, France

1- Objectives

Forage protein value evaluation relies on the estimation of forage protein degradability in the rumen (deg).

Deg is affected by botanical family, plant species, growth cycle and level of nitrogen fertilization, and decreases with plant maturity stage.

→ To develop equations for predicting forage deg values in order to improve the estimation of forage protein values expressed in PDI.

3 - Results

Mean values in the data base for fresh forages

		n	deg (% of CP)	CP (g kg ⁻¹ DM)
Grasses	1 st growth	60	80.9 ± 6.5	150 ± 52
	Other growths	141	73.9 ± 8.5	197 ± 61
Legumes	1 st growth	43	81.4 ± 4.9	185 ± 46
	Other growths	33	81.4 ± 4.8	205 ± 49
Natural grasslands	1 st growth	31	73.6 ± 9.0	158 ± 54
	Other growths	67	67.2 ± 8.3	194 ± 42

• Fresh forages

$$\text{deg} = 51.2 + 0.14 \text{ CP} - 0.00017 \text{ CP}^2$$

$$R^2 = 0.87; \text{RSD} = 3.5$$


- + 8.8 (grasses - 1st growth)
- + 4.6 (grasses - other growths)
- + 6.8 (legume forages - all growths)
- + 4.4 (natural grasslands - 1st growth)
- + 0 (natural grasslands - other growths)

• Conserved forages

$$\text{deg}_{\text{silage}} = 46.5 + 0.56 \text{ deg}_{\text{fresh forage}} - 0.25 \text{ DM}_{\text{silage}} \quad (R^2 = 0.53, \text{RSD} = 4.2)$$

$$\text{deg}_{\text{hay}} = 17.1 + 0.68 \text{ deg}_{\text{fresh forage}} \quad (R^2 = 0.71, \text{RSD} = 3.6)$$

• Example for natural grasslands – 1st growth



2 - Methods

Fresh forages : meta-analysis of a data-base of 375 *in situ* deg measurements for fresh forages. Data from :


INRA : 11 trials 214 data

European literature and personal communications : 12 trials 161 data

Conserved forages : for the same forages comparison between

- Fresh and hay (29 data)
- Fresh and silages (32 data)

deg – CP relationship for the 23 trials



4 - Conclusion

These equations have satisfactory accuracy and will be used to improve estimation for herbage from European temperate climate areas in the INRA PDI system.